



(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number  
**WO 03/086306 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K**
- (21) International Application Number: PCT/US03/11076
- (22) International Filing Date: 11 April 2003 (11.04.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
 60/372,058 12 April 2002 (12.04.2002) US  
 60/405,495 23 August 2002 (23.08.2002) US  
 60/434,607 18 December 2002 (18.12.2002) US  
 10/410,991 10 April 2003 (10.04.2003) US
- (71) Applicant (for all designated States except US): **ARENA PHARMACEUTICALS, INC.** [US/US]; 6166 Nancy Ridge Drive, San Diego, CA 92121 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SMITH, Jeffrey** [US/US]; 8041 Jade Coast Road, San Diego, CA 92126 (US). **SMITH, Brian** [US/US]; 11184 Vista Sorrento Parkway, Apartment G-301, San Diego, CA 92130 (US).
- (74) Agents: **STRAHER, Michael, P.** et al.; Cozen O'Connor, 1900 Market Street, Philadelphia, PA 19103 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
 — without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 5HT<sub>2C</sub> RECEPTOR MODULATORS

(57) Abstract: The present invention relates to novel compounds of: (Formula I); which act as 5HT<sub>2C</sub> receptor modulators. These compounds are useful in pharmaceutical compositions whose use includes the treatment of obesity.

## 5HT<sub>2C</sub> RECEPTOR MODULATORS

### Cross-Reference to Related Applications

This application claims priority benefit of U.S. Provisional Application Number 60/372,058, filed April 12, 2002; U.S. Provisional Patent Application  
5 Number 60/405,495, filed August 23, 2002, U.S. Provisional Patent Application Number 60/434,607, filed December 18, 2002, and U.S. Non-Provisional Patent Application No. (not yet assigned), filed April 10, 2003, which are hereby incorporated by reference in their entirety.

### Field of the Invention

10 The present invention relates to compounds which act as modulators of 5HT<sub>2C</sub> receptors, compositions including the compounds, and methods of using the compounds.

### Background of the Invention

Obesity is a life-threatening disorder in which there is an increased risk of  
15 morbidity and mortality arising from concomitant diseases such as type II diabetes, hypertension, stroke, cancer and gallbladder disease.

Obesity is now a major healthcare issue in the Western World and increasingly in some third world countries. The increase in numbers of obese people is due largely to the increasing preference for high fat content foods but  
20 also, and this can be a more important factor, the decrease in activity in most people's lives. In the last 10 years there has been a 30% increase in the incidence of obesity in the USA and that about 30% of the population of the USA is now considered obese.

Whether someone is classified as overweight or obese is generally  
25 determined on the basis of their body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m<sup>2</sup>). Thus, the units of BMI are kg/m<sup>2</sup> and it is possible to calculate the BMI range associated with minimum mortality in each decade of life. Overweight is defined as a BMI in the range 25-30 kg/m<sup>2</sup>, and obesity as a BMI greater than 30 kg/m<sup>2</sup> (see TABLE below).

**CLASSIFICATION OF WEIGHT BY  
BODY MASS INDEX (BMI)**

BMI	CLASSIFICATION
< 18.5	Underweight
18.5-24.9	Normal
25.0-29.9	Overweight
30.0-34.9	Obesity (Class I)
35.0-39.9	Obesity (Class II)
>40	Extreme Obesity (Class III)

As the BMI increases there is an increased risk of death from a variety of causes that is independent of other risk factors. The most common diseases with obesity are cardiovascular disease (particularly hypertension), diabetes (obesity aggravates the development of diabetes), gall bladder disease (particularly cancer) and diseases of reproduction. Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk of developing coronary heart disease.

There are problems however with the BMI definition in that it does not take into account the proportion of body mass that is muscle in relation to fat (adipose tissue). To account for this, obesity can also be defined on the basis of body fat content: greater than 25% in males and 30% in females.

Obesity considerably increases the risk of developing cardiovascular diseases as well. Coronary insufficiency, atheromatous disease, and cardiac insufficiency are at the forefront of the cardiovascular complication induced by obesity. It is estimated that if the entire population had an ideal weight, the risk of coronary insufficiency would decrease by 25% and the risk of cardiac insufficiency and of cerebral vascular accidents by 35%. The incidence of coronary diseases is doubled in subjects less than 50 years of age who are 30% overweight. The diabetes patient faces a 30% reduced lifespan. After age 45, people with diabetes are about three times more likely than people without diabetes to have significant heart disease and up to five times more likely to have a stroke. These findings emphasize the inter-

relations between risks factors for NIDDM and coronary heart disease and the potential value of an integrated approach to the prevention of these conditions based on the prevention of obesity (Perry, I. J., et al., *BMJ* 310, 560-564 (1995)).

Diabetes has also been implicated in the development of kidney disease, eye diseases and nervous-system problems. Kidney disease, also called nephropathy, occurs when the kidney's "filter mechanism" is damaged and protein leaks into urine in excessive amounts and eventually the kidney fails. Diabetes is also a leading cause of damage to the retina at the back of the eye and increases risk of cataracts and glaucoma. Finally, diabetes is associated with nerve damage, especially in the legs and feet, which interferes with the ability to sense pain and contributes to serious infections. Taken together, diabetes complications are one of the nation's leading causes of death.

The first line of treatment is to offer diet and life style advice to patients such as reducing the fat content of their diet and increasing their physical activity. However many patients find this difficult and need additional help from drug therapy to maintain results from these efforts.

Most currently marketed products have been unsuccessful as treatments for obesity owing to a lack of efficacy or unacceptable side-effect profiles. The most successful drug so far was the indirectly acting 5-hydroxytryptamine (5-HT) agonist d-fenfluramine (Redux<sup>TM</sup>) but reports of cardiac valve defects in up to one third of patients led to its withdrawal by the FDA in 1998.

In addition, two drugs have recently been launched in the USA and Europe: Orlistat (Xenical<sup>TM</sup>), a drug that prevents absorption of fat by the inhibition of pancreatic lipase, and Sibutramine (Reductil<sup>TM</sup>), a 5-HT/noradrenaline re-uptake inhibitor. However, side effects associated with these products may limit their long-term utility. Treatment with Xenical<sup>TM</sup> is reported to induce gastrointestinal distress in some patients, while Sibutramine has been associated with raised blood pressure in some patients.

Serotonin (5-HT) neurotransmission plays an important role in numerous physiological processes both in health and in psychiatric disorders. 5-HT has been implicated in the regulation of feeding behavior for some time. 5-HT appears to




work by inducing a feeling of fullness or satiety so eating stops earlier and fewer calories are consumed. It has been shown that a stimulatory action of 5-HT on the 5HT<sub>2C</sub> receptor plays an important role in the control of eating and in the anti-obesity effect of d-fenfluramine. As the 5-HT<sub>2C</sub> receptor is expressed in high density in the brain (notably in the limbic structures, extrapyramidal pathways, thalamus and hypothalamus i.e. PVN and DMH, and predominantly in the choroid plexus) and is expressed in low density or is absent in peripheral tissues, a selective 5-HT<sub>2C</sub> receptor agonist can be a more effective and safe anti-obesity agent. Also, 5-HT<sub>2C</sub> knockout mice are overweight with cognitive impairment and susceptibility to seizure.

It is believed that 5HT<sub>2C</sub> may play a role in obsessive compulsive disorder, some forms of depression, and epilepsy. Accordingly, agonists can have anti-panic properties, and properties useful for the treatment of sexual dysfunction.

In sum, the 5HT<sub>2C</sub> receptor is a validated and well-accepted receptor target for the treatment of obesity and psychiatric disorders, and it can be seen that there is a need for selective 5HT<sub>2C</sub> agonists which safely decrease food intake and body weight. The present invention is directed to these, as well as other, important ends.

### Summary of the Invention

The present invention, in one aspect, relates to compounds represented by Formula (I):



wherein:

R<sub>1</sub> is H or C<sub>1-8</sub> alkyl;

R<sub>2</sub> is C<sub>1-8</sub> alkyl, -CH<sub>2</sub>-O-C<sub>1-8</sub> alkyl, -C(=O)-O-C<sub>1-8</sub> alkyl, -C(=O)-NH-C<sub>1-8</sub> alkyl, or CH<sub>2</sub>OH;

R<sub>2a</sub> is H or CH<sub>3</sub>;

or R<sub>2</sub> and R<sub>2a</sub> together form -CH<sub>2</sub>-CH<sub>2</sub>-;

R<sub>3</sub> and R<sub>4</sub> are each independently H, halogen, perhalo alkyl, (preferably CF<sub>3</sub>), CN, OR<sub>5</sub>, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl; and

R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(A) if R<sub>2</sub> is methyl and R<sub>1</sub> and R<sub>3</sub> are both H, then R<sub>4</sub> is not thiazole, substituted thiazole or a thiazole derivative:

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H;

(C) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be NHR<sub>5</sub> or N(R<sub>5</sub>)<sub>2</sub>;

(D) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be imidazole, substituted imidazole, or an imidazole derivative; and

(E) if R<sub>3</sub> is OH, and R<sub>1</sub> is methyl then R<sub>2</sub> cannot be cyclopentyl, -CH<sub>2</sub>-cyclohexyl, cyclopropylmethyl, or cyclohexyl.

In some embodiments of the compounds and methods of the invention, when R<sub>1</sub>, R<sub>2a</sub>, R<sub>3</sub> and R<sub>6</sub> are H and R<sub>2</sub> is methyl, then R<sub>4</sub> cannot be a chlorine atom.

In other embodiments of the compounds and methods of the invention, when R<sub>1</sub>, R<sub>2a</sub>, R<sub>3</sub> and R<sub>6</sub> are H and R<sub>2</sub> is methyl, then R<sub>4</sub> can be a chlorine atom.

In some alternate embodiments of the compounds of Formula (I), if R<sub>4</sub> is OR<sub>5</sub>, then R<sub>2</sub> cannot be alkyl.

In some embodiments of the compounds of Formula (I), R<sub>1</sub> is H. In some embodiments of the compounds of Formula (I), R<sub>1</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>1</sub> is methyl. In some embodiments of the compounds of Formula (I), R<sub>1</sub> is n-propyl.

5            In some embodiments of the compounds of Formula (I), R<sub>2</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>2</sub> is methyl. In some embodiments of the compounds of Formula (I), R<sub>2</sub> is ethyl. In some embodiments of the compounds of Formula (I), R<sub>2</sub> is isopropyl. In some embodiments of the compounds of Formula (I), R<sub>2</sub> and R<sub>2a</sub> together form -CH<sub>2</sub>-CH<sub>2</sub>-.

10           In some embodiments of the compounds of Formula (I), R<sub>3</sub> is halogen. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is chlorine. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is bromine. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is iodine. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is perhaloalkyl. In some  
15           embodiments of the compounds of Formula (I), R<sub>3</sub> is CF<sub>3</sub>. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S. In some embodiments, R<sub>3</sub> is a radical derived from thiophenyl, furanyl, pyrrolyl, pyrazolyl or imidazolyl.

             In some embodiments of the compounds of Formula (I), R<sub>4</sub> is perhaloalkyl.  
20           In some embodiments of the compounds of Formula (I), R<sub>4</sub> is CF<sub>3</sub>. In some embodiments of the compounds of Formula (I), R<sub>4</sub> is -OR<sub>5</sub>. In some embodiments R<sub>5</sub> is methyl, ethyl, n-propyl, isopropyl or allyl. In some embodiments of the compounds of Formula (I), R<sub>5</sub> is methyl or allyl. In some  
25           embodiments of the compounds of Formula (I), R<sub>4</sub> is a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S, and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl. In some embodiments, R<sub>4</sub> is a radical derived from thiophenyl, furanyl, pyrrolyl, pyrazolyl or imidazolyl, which can optionally be mono- or di-substituted selected from halogen or methyl. In  
30           some embodiments of the compounds of Formula (I), R<sub>4</sub> is phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and

alkoxy. In some embodiments of the compounds of Formula (I), R<sub>3</sub> and R<sub>4</sub> taken together form -O-CH=C(CH<sub>3</sub>)-.

In some embodiments of the compounds of Formula (I), R<sub>3</sub> is halogen and R<sub>4</sub> is

5 -OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is chlorine and R<sub>4</sub> is -OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is bromine and R<sub>4</sub> is -OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is iodine and R<sub>4</sub> is -OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the  
10 compounds of Formula (I), R<sub>3</sub> is halogen and R<sub>4</sub> is methoxy. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is halogen and R<sub>4</sub> is allyloxy.

In some embodiments of the compounds of Formula (I), R<sub>3</sub> is H and R<sub>4</sub> is a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S, and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, or R<sub>4</sub> is  
15 phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy.

In some embodiments of the compounds of Formula (I), R<sub>3</sub> is H and R<sub>4</sub> is a disubstituted pyrrazole or monohalo-substituted phenyl. In some such  
20 embodiments of the compounds of Formula (I), the substituents of the pyrrazole are bromine and methyl.

In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub>. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is aryl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub>  
25 wherein R<sub>5</sub> is heteroaryl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is arylalkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is arylmethyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is heteroarylalkyl. In some  
30 embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is heteroarylalkylmethyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is

OR<sub>5</sub> wherein R<sub>5</sub> is perhaloalkyl. In some embodiments of the compounds of Formula (I), R<sub>3</sub> is OR<sub>5</sub> wherein R<sub>5</sub> is allyl.

In some embodiments of the compounds of Formula (I):

5 R<sub>2</sub> is methyl, ethyl, isopropyl, or CH<sub>2</sub>OH; or R<sub>2</sub> and R<sub>2a</sub> taken together form -CH<sub>2</sub>-CH<sub>2</sub>-;

R<sub>3</sub> is H, halogen, or a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S, and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

10 R<sub>4</sub> is H, alkoxy, a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, or phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy;

or R<sub>3</sub> and R<sub>4</sub> taken together form -O-CH=C(CH<sub>3</sub>)-; and

15 R<sub>6</sub> is H or methyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In some embodiments of the compounds of Formula (I):

R<sub>2</sub> is methyl, ethyl, isopropyl, or CH<sub>2</sub>OH; or R<sub>2</sub> and R<sub>2a</sub> taken together form -CH<sub>2</sub>-CH<sub>2</sub>-;

R<sub>3</sub> is chlorine, bromine, or iodine;

20 R<sub>4</sub> is alkoxy; and

R<sub>6</sub> is H or methyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In some embodiments of the compounds of Formula (I):

R<sub>1</sub> is H;

25 R<sub>2</sub> is methyl;

R<sub>3</sub> is H, chlorine, bromine, or thiophene;

R<sub>4</sub> is alkoxy, pyrrazoly-3-yl or phenyl wherein said pyrrazole optionally has up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, and said phenyl optionally has a single halogen substituent; and

30

R<sub>6</sub> is H; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In some embodiments of the compounds of Formula (I), the compound is a member of the group consisting of: 8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Benzoyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Propyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 3,5-Dimethyl-6,7,8,9-tetrahydro-5*H*-1-oxa-7-aza-cycloheptaindene; 7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-methoxy-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(4-Bromo-2-methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 1-Methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Propyl-8-

iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 1-Ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(4-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 8-bromo-1-methoxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In some embodiments of the compounds of Formula (I), the compound is a member of the group consisting of: 8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In some embodiments of the compounds of Formula (I), the compound is a member of the group consisting of: 8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-7-fluoro-1-

methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 8-Chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

5 The present invention also provides compositions comprising one or more compounds of the invention, and one or more pharmaceutically acceptable carriers.

The present invention further provides methods of modulating a 5HT<sub>2C</sub> receptor comprising contacting said receptor with a pharmaceutically effective amount of a compound or composition of the invention. Preferably, said  
10 compound is an agonist of said receptor.

The present invention further provides methods of prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea comprising administering to a patient in need of such  
15 prophylaxis or treatment an effective dose of a compound of the invention.

In some embodiments, the disorders of the central nervous system include depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated  
20 with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related mental disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension. In some  
25 embodiments, the disorders of the central nervous system is obesity.

In some embodiments, the damage to the central nervous system is by trauma, stroke, neurodegenerative diseases or toxic or infective CNS diseases, including encephalitis and meningitis.

In some embodiments, the cardiovascular disorder thrombosis. In further  
30 embodiments, the gastrointestinal disorder is dysfunction of gastrointestinal motility.



The present invention further provides methods of decreasing food intake of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound or composition of the invention.

5 The present invention further provides methods of inducing satiety in a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound or composition of the invention.

The present invention further provides methods of controlling weight gain of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound or composition of the invention.

10 The present invention further provides methods of treating obesity comprising administering to a patient in need of such treatment a pharmaceutically effective amount of a compound or composition of the invention.

In some embodiments, some of the foregoing methods of the invention further comprising the step of identifying a subject, said subject being in need of decreasing food intake, controlling weight gain, or treating obesity, wherein said identifying step is performed prior to administering to said subject said pharmaceutically effective amount of said compound or composition of the invention.

20 One aspect of the present invention pertains to a compound of Formula (I) for use in a method of treatment of the human or animal body by therapy.

One aspect of the present invention pertains to a compound of Formula (I) for use in a method of prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea. In some  
25 embodiments the disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy,  
30 personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood,

aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension. In some embodiments the disorder is obesity.

One aspect of the present invention pertains to a compound of Formula (I) for the manufacture of a medicament for use in the prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea. In some embodiments the disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension. In some embodiments the disorder is obesity.

In some embodiments, the invention provides methods for alleviation of a symptom of any of the diseases, conditions or disorders mentioned herein.

Applicants reserve the right to exclude any one or more of the compounds from any of the embodiments of the invention. Applicant additionally reserves the right to exclude any disorder from any of the embodiments of the invention.

#### **Brief Description of the Figures**

Figures 1A-1G illustrate the effects of seven different compounds of the invention on food intake in food-deprived rats.


#### **Detailed Description of the Invention**

The present invention relates to 5HT<sub>2C</sub> receptor agonist compounds, methods of modulating 5HT<sub>2C</sub> receptors by contacting the receptors with one or more compounds of the invention. The present invention also relates to methods of decreasing food intake, controlling weight gain, or treating obesity, using compounds of the invention.

The term "antagonist" is intended to mean moieties that competitively bind to the receptor at the same site as agonists (for example, the endogenous ligand), but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists or partial agonists. Antagonists do not diminish the baseline intracellular response in the absence of an agonist or partial agonist. As used herein, the term "agonist" is intended to mean moieties that activate the intracellular response when they bind to the receptor, or enhance GTP binding to membranes. In the context of the present invention, a pharmaceutical composition comprising a 5HT<sub>2C</sub> receptor agonist of the invention can be utilized for modulating the activity of the 5HT<sub>2C</sub> receptor, decreasing food intake, inducing satiation (i.e., the feeling of fullness), controlling weight gain, treating obesity, decreasing body weight and/or affecting metabolism such that the recipient loses weight and/or maintains weight. Such pharmaceutical compositions can be used in the context of disorders and/or diseases where weight gain is a component of the disease and/or disorder such as, for example, obesity.

As used herein, the term "contact" or "contacting" shall mean bringing the indicated moieties together, whether in an in vitro system or an in vivo system. Thus, "contacting" an 5HT<sub>2C</sub> receptor with a compound of the invention includes the administration of a compound of the invention to an animal having an 5HT<sub>2C</sub> receptor, as well as, for example, introducing a compound of the invention into a sample containing a cellular or more purified preparation containing an 5HT<sub>2C</sub> receptor.

Compounds of the invention include those having the Formula (I), shown below:



wherein:

R<sub>1</sub> is H or C<sub>1-8</sub> alkyl;

R<sub>2</sub> is C<sub>1-8</sub> alkyl, -CH<sub>2</sub>-O-C<sub>1-8</sub> alkyl, -C(=O)-O-C<sub>1-8</sub> alkyl, -C(=O)-NH-C<sub>1-8</sub> alkyl, or CH<sub>2</sub>OH;

R<sub>2a</sub> is H; or R<sub>2</sub> and R<sub>2a</sub> together form -CH<sub>2</sub>-CH<sub>2</sub>-;

5 R<sub>3</sub> and R<sub>4</sub> are each independently H, halogen, perhalo alkyl, (preferably CF<sub>3</sub>), CN, OR<sub>5</sub>, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

10 or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl; and

15 R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(A) if R<sub>2</sub> is methyl and R<sub>1</sub> and R<sub>3</sub> are both H, then R<sub>4</sub> is not thiazole, substituted thiazole or a thiazole derivative:

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H;

20 (C) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be NHR<sub>5</sub> or N(R<sub>5</sub>)<sub>2</sub>;

(D) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be imidazole, substituted imidazole, or an imidazole derivative; and

25 (E) if R<sub>3</sub> is OH, and R<sub>1</sub> is methyl then R<sub>2</sub> cannot be cyclopentyl, -CH<sub>2</sub>-cyclohexyl, cyclopropylmethyl, or cyclohexyl;

or provided (A), (B), (C), (D) above, and if R<sub>4</sub> is OR<sub>5</sub>, then R<sub>2</sub> cannot be alkyl.

30 In some embodiments of the compounds and methods of the invention, when R<sub>1</sub>, R<sub>2a</sub>, R<sub>3</sub> and R<sub>6</sub> are H and R<sub>2</sub> is methyl, then R<sub>4</sub> cannot be a chlorine atom.

In other embodiments of the compounds and methods of the invention, when  $R_1$ ,  $R_{2a}$ ,  $R_3$  and  $R_6$  are H and  $R_2$  is methyl, then  $R_4$  can be a chlorine atom.

5 In some embodiments, if  $R_4$  is  $OR_5$ , then  $R_2$  cannot be cyclopentyl,  $-CH_2-$ cyclohexyl, 3,3-dimethyl-2-allyl, 3,3-dimethyl-2-methylallyl, 2-methylallyl, 2-butenyl, cyclopropylmethyl, cyclohexyl, or allyl.

10 It will be appreciated that compounds of Formula (I) may have one or more chiral centers, and exist as enantiomers or diastereomers. The invention is to be understood to extend to all such enantiomers, diastereomers and mixtures thereof, including racemates. Formula (I) and the formulae hereinafter are intended to represent all individual isomers and mixtures thereof, unless stated or shown otherwise.

15 As used herein, the term "alkyl" is intended to denote hydrocarbon groups including straight chain, branched and cyclic hydrocarbons, including for example but not limited to methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, sec-butyl, tert-butyl, cyclobutyl, cyclopropylmethyl, n-pentyl, isopentyl, tert-pentyl, cyclopentyl, cyclopentylmethyl, n-hexyl, cyclohexyl, and the like. Throughout this specification, it should be understood that the term alkyl is intended to encompass both non-cyclic hydrocarbon groups and cyclic hydrocarbon groups.

20 In some embodiments of the compounds of the invention, alkyl groups are non-cyclic. In further embodiments, alkyl groups are cyclic, and in further embodiments, alkyl groups are both cyclic and noncyclic. Where no preference is specified, the term "alkyl" is intended to denote groups that are both cyclic and non-cyclic.

25 As used herein, the term "alkenyl" is intended to denote hydrocarbon compounds including straight chain, branched and cyclic hydrocarbons that contain at least one double bond, including for example but not limited to allyl, 2-methyl-allyl, 4-but-3-enyl, 4-hex-5-enyl, 3-methyl-but-2-enyl, cyclohex-2-enyl and the like.

30 As used herein, the term "halogen" has its normal meaning of period seven elements, including F, Cl, Br and I.

The term "alkoxy" is intended to denote substituents of the formula -O-alkyl, including -O-allyl. The term "lower" when used in connection with substituents such as alkyl indicates 6 carbons or less.

5 The term "arylalkyl" or "aralkyl" is intended to denote an alkyl group that bears an aryl substituent, for example a benzyl group. The term "alkylaryl" or "alkaryl" is intended to denote an aryl group that bears an alkyl substituent, for example a 4-methylphenyl group.

10 As used herein, the term "aryl" is intended to mean monocyclic and polycyclic aromatic groups. Although aryl groups can include as few as 3 carbon atoms, preferred aryl groups have 6 to about 14 carbon atoms, more preferably 6 to about 10 carbon atoms. Examples of aryl groups include but are not limited to phenyl, naphthyl, anthracyl, phenanthryl and pyrenyl.

15 The term "heteroaryl" is intended to denote an aryl group that contains at least one, and preferably from one to four ring "hetero" (i.e., non-carbon, e.g., O, N or S) atom. Examples of "heteroaryl" groups are radicals derived from 5- and 6-member aryl ring compounds having from one to four nitrogen, sulfur and/or oxygen atoms, for example pyrrole, pyrazole, imidazole, triazole, tetrazole, pyridine, pyrimidine, furan, pyran, thiophene, benzimidazole, quinoline, isoquinoline, oxazole, thiazole, and thiadiazole.

20 As used herein the term heteroarylalkyl means an alkyl group that bears a heteroaryl substituent, for example a group having the structure -CH<sub>2</sub>-pyrrole-2-yl.

25 The term "substituted thiazole" means a radical derived from thiazole that bears at least one substituent group. The term "thiazole derivative" means a fused ring system in which one of the fused rings is thiazole.

The term "substituted imidazole" means a radical derived from imidazole that bears at least one substituent group. The term "imidazole derivative" means a fused ring system in which one of the fused rings is imidazole.

30 In some embodiments of the invention, R<sub>4</sub> is OR<sub>5</sub>. In some such embodiments, R<sub>2</sub> cannot be cyclopentyl, -CH<sub>2</sub>-cyclohexyl, 3,3-dimethyl-2-allyl, 3,3-dimethyl-2-methylallyl, 2-methylallyl, 2-butenyl, cyclopropylmethyl,

cyclohexyl, or allyl. In further such embodiments, R<sub>2</sub> cannot be alkyl.

In some embodiments of the compounds of Formula (I), R<sub>3</sub> is halogen and R<sub>4</sub> is

5 -OR<sub>5</sub>. In some embodiments of the compounds of Formula (I), R<sub>5</sub> is allyl, 2-methyl-allyl, 4-but-3-enyl, 4-hex-5-enyl, 3-methyl-but-2-enyl or cyclohex-2-enyl. In some embodiments of the compounds of Formula (I), R<sub>5</sub> is methyl, ethyl, n-propyl, isopropyl or allyl. In some embodiments of the compounds of Formula (I), R<sub>5</sub> is methyl or allyl.

10 Certain substituents of the compounds disclosed herein can optionally be substituted, i.e., they can optionally bear further substituent groups. Some preferred substituent groups include halogen, lower alkyl (including but not limited to methyl, ethyl, isopropyl, cyclopropyl, tert-butyl, and methylcyclopropyl), alkoxy, mono-, di- or trihaloalkoxy (e.g., -O-CX<sub>3</sub> where X is halogen), -(CH<sub>2</sub>)<sub>y</sub>NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>y</sub>NHBoc,  
15 -N(R<sub>4a</sub>)(R<sub>4b</sub>), phenyl, methoxyphenyl and naphthyl.

At various places in the present specification substituents of compounds of the invention are disclosed in groups or in ranges. It is specifically intended that the invention include each and every individual subcombination of the members of such groups and ranges. For example, the term "C<sub>1-8</sub> alkyl" is specifically  
20 intended to individually disclose methyl, ethyl, C<sub>3</sub> alkyl, C<sub>4</sub> alkyl, C<sub>5</sub> alkyl, C<sub>6</sub> alkyl, C<sub>7</sub> alkyl and C<sub>8</sub> alkyl.

In a preferred embodiment, the compounds of Formula (I) are selected from:

25 8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Benzyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Propyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Hydroxy-8-iodo-1-methyl-2,3,4,5-  
30 tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 3,5-Dimethyl-6,7,8,9-tetrahydro-5*H*-1-oxa-7-aza-

cycloheptaindene; 7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-7-methoxy-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(4-Bromo-2-methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 1-Methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Propyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 1-Ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(2-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(3-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-(4-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 8-bromo-1-methoxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.



In a preferred embodiment, the compounds of Formula (I) are selected from:

N-methyl-8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Iodo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and N-methyl-7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In a preferred embodiment, the compounds of Formula (I) are selected from:

N-methyl-8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-7,8-Dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-methyl-8-Chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and N-methyl-8-Chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In a preferred embodiment, the compounds of Formula (I) are selected from:

8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-7-

methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

In a preferred embodiment, the compounds of Formula (I) are selected from:

8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 7,8-Dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; 8-Chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and 8-Chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

The compounds of the invention may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. The compounds can be, for example, racemates or optically active forms. The optically active forms can be obtained by resolution of the racemates or by asymmetric synthesis. In some embodiments, compounds of Formula (I) are *R* enantiomers. In some embodiments, compounds of Formula (I) are *S* enantiomers. In some embodiments, compounds of Formula (I) are varying mixtures of enantiomers.

According to a further aspect of the invention, compounds of Formula (I) are provided for use in therapy. The compounds of Formula (I) can be used in the

prophylaxis or treatment of disorders associated with 5-HT<sub>2C</sub> receptor function.

The compounds of Formula (I) can be used in the prophylaxis or treatment of central nervous disorders such as depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or  
5 panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory  
10 disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa or premenstrual tension; damage of the central nervous system such as by trauma, stroke, neurodegenerative diseases or toxic or infective CNS diseases such as encephalitis or meningitis; cardiovascular disorders such as thrombosis; gastrointestinal disorders such as dysfunction of gastrointestinal  
15 motility; diabetes insipidus; and sleep apnea.

According to a further aspect of the invention, there is provided use of a compound of Formula (I) in the manufacture of a medicament for the prophylaxis or treatment of the disorders disclosed herein. In a preferred embodiment, there is provided a use of a compound of Formula (I) in the manufacture of a medicament  
20 for the prophylaxis or treatment of obesity.

The compounds according to the invention may optionally exist as pharmaceutically acceptable salts including pharmaceutically acceptable acid addition salts prepared from pharmaceutically acceptable non-toxic acids including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, dichloroacetic, formic, fumaric,  
25 gluconic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, oxalic, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, oxalic, p-toluenesulfonic and the like, such as the pharmaceutically acceptable salts listed in Journal of Pharmaceutical  
30 Science, 66, 2 (1977) and incorporated herein by reference.

The acid addition salts can be obtained as the direct products of compound

synthesis. In the alternative, the free base can be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent. The compounds of this invention may form solvates with standard low molecular weight solvents using methods known to the skilled artisan.

Compositions of the invention may conveniently be administered in unit dosage form and can be prepared by any of the methods well known in the pharmaceutical art, for example, as described in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., Easton, PA, 1980).

The compounds of the invention can be employed as the sole active agent in a pharmaceutical or can be used in combination with other active ingredients which could facilitate the therapeutic effect of the compound.

Compounds of the present invention or a solvate or physiologically functional derivative thereof can be used as active ingredients in pharmaceutical compositions, specifically as 5HT<sub>2C</sub> receptor agonists. By the term "active ingredient" is defined in the context of a "pharmaceutical composition" and shall mean a component of a pharmaceutical composition that provides the primary pharmaceutical benefit, as opposed to an "inactive ingredient" which would generally be recognized as providing no pharmaceutical benefit. The term "pharmaceutical composition" shall mean a composition comprising at least one active ingredient and at least one ingredient that is not an active ingredient (for example and not limitation, a filler, dye, or a mechanism for slow release), whereby the composition is amenable to use for a specified, efficacious outcome in a mammal (for example, and not limitation, a human).

The data developed herein supports the conclusion that the presently disclosed 5HT<sub>2C</sub> receptor agonists are of use for the treatment or prophylaxis of clinical obesity or overweight disorders in mammals, including, but not limited to, human. Compounds of the present invention can be administered by oral, sublingual, parenteral, rectal, topical administration or by a transdermal patch. Transdermal patches dispense a drug at a controlled rate by presenting the drug for absorption in an efficient manner with a minimum of degradation of the drug.

Typically, transdermal patches comprise an impermeable backing layer, a single pressure sensitive adhesive and a removable protective layer with a release liner. One of ordinary skill in the art will understand and appreciate the techniques appropriate for manufacturing a desired efficacious transdermal patch based upon the needs of the artisan.

In addition to the neutral forms of compounds of the present invention, by appropriate addition of an ionizable substituent, which does not alter the receptor specificity of the compound, physiologically acceptable salts of the compounds may also be formed and used as therapeutic agents. Different amounts of the compounds of the present invention will be required to achieve the desired biological effect. The amount will depend on factors such as the specific compound, the use for which it is intended, the means of administration, and the condition of the treated individual – all of these dosing parameters are within the level of one of ordinary skill in the medicinal arts. A typical dose can be expected to fall in the range of 0.001 to 200 mg per kilogram of body weight of the mammal. Unit doses may contain from 1 to 200 mg of the compounds of the present invention and can be administered one or more times a day, individually or in multiples.

Pharmaceutical compositions, including, but not limited to, pharmaceutical compositions, comprising at least one compound of the present invention and/or an acceptable salt or solvate thereof (*e.g.*, a pharmaceutically acceptable salt or solvate) as an active ingredient combined with at least one carrier or excipient (*e.g.*, pharmaceutical carrier or excipient) can be used in the treatment of clinical conditions for which a 5HT<sub>2C</sub> receptor agonist is indicated. At least one compound of the present invention can be combined with the carrier in either solid or liquid form in a unit dose formulation. The pharmaceutical carrier must be compatible with the other ingredients in the composition and must be tolerated by the individual recipient. Other physiologically active ingredients can be incorporated into the pharmaceutical composition of the invention if desired, and if such ingredients are compatible with the other ingredients in the composition. Formulations can be prepared by any suitable method, typically by uniformly

mixing the active compound(s) with liquids or finely divided solid carriers, or both, in the required proportions, and then, if necessary, forming the resulting mixture into a desired shape.

5           Conventional excipients, such as binding agents, fillers, acceptable wetting agents, tableting lubricants, and disintegrants can be used in tablets and capsules for oral administration. Liquid preparations for oral administration can be in the form of solutions, emulsions, aqueous or oily suspensions, and syrups. Alternatively, the oral preparations can be in the form of dry powder that can be reconstituted with water or another suitable liquid vehicle before use. Additional  
10           additives such as suspending or emulsifying agents, non-aqueous vehicles (including edible oils), preservatives, and flavorings and colorants can be added to the liquid preparations. Parenteral dosage forms can be prepared by dissolving the compound of the invention in a suitable liquid vehicle and filter sterilizing the solution before filling and sealing an appropriate vial or ampoule. These are just a  
15           few examples of the many appropriate methods well known in the art for preparing dosage forms.


          It is noted that when the 5HT<sub>2C</sub> receptor agonists are utilized as active ingredients in a pharmaceutical composition, these are not intended for use only in humans, but in other non-human mammals as well. Indeed, recent advances in the  
20           area of animal health-care mandate that consideration be given for the use of 5HT<sub>2C</sub> receptor agonists for the treatment of obesity in domestic animals (*e.g.*, cats and dogs), and 5HT<sub>2C</sub> receptor agonists in other domestic animals where no disease or disorder is evident (*e.g.*, food-oriented animals such as cows, chickens, fish, etc.). Those of ordinary skill in the art are readily credited with  
25           understanding the utility of such compounds in such settings.

          The compounds of the present invention can be readily prepared according to a variety of synthetic manipulations, all of which would be familiar to one skilled in the art. A representative general synthesis is set forth below in Scheme  
I:

## Scheme I

## GENERAL REACTION SCHEME

5




Those of skill in the art will appreciate that a wide variety of compounds of the invention can be prepared according to Scheme I. For example, by starting with an appropriately substituted 2-phenyl ethylamino compound A having any of a wide variety of substituents R<sub>1</sub> and R<sub>2</sub>, the corresponding 7- and/or 8-substituted  
5 1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (compound H) can be prepared.

N-alkylation can be accomplished by, for example, treatment with excess paraformaldehyde (for methylation) or a higher order aldehyde, followed by reduction with NaBH<sub>3</sub>CN according to the general procedure of synthetic examples 9 and 10, *infra*. In addition, by starting with an appropriately substituted  
10 1-alkyl-2-phenyl ethylamino compound A having any of a wide variety of substituents R<sub>1</sub> and R<sub>2</sub>, the corresponding 7- and/or 8-substituted 2,5-dialkyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine compound can be prepared.

In the synthesis of many compounds of the invention, protecting groups can be required to protect various functionality or functionalities during the  
15 synthesis. Representative protecting groups suitable for a wide variety of synthetic transformations are disclosed in Greene and Wuts, *Protective Groups in Organic Synthesis*, 2d ed, John Wiley & Sons, New York, 1991, the disclosure of which is incorporated herein by reference in its entirety.

As will be recognized, the steps of the methods of the present invention  
20 need not be performed any particular number of times or in any particular sequence. Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples thereof, which are intended to be illustrative and not intended to be limiting.



**Examples****Synthetic Examples****Example 1: (R,S) 8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine***N-Trifluoroacetyl-3-methoxyphenethylamine*

A solution of 3-methoxyphenethylamine (10.0 g, 64.0 mmol) in dichloromethane (150 mL), was cooled to 0 C, and treated with pyridine (6.5 mL, 83.5 mmol) followed by the dropwise addition of trifluoroacetic anhydride (17.9 g, 83.5 mmol) and the resulting mixture stirred for 3 hours while warming to 20 C. The product mixture was diluted with EtOAc (500 mL), washed sequentially with 10% aqueous HCl (100 mL), water (100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 15.8 g of a yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.26 (dd, J=8, 8 Hz, 1 H), 6.81 (d, J=8 Hz, 1 H), 6.77 (d, J=8 Hz, 1 H), 6.72 (s, 1 H), 6.30 (bs, 1 H), 3.80 (s, 3 H), 3.62 (dd, J=7, 7 Hz, 2 H), 2.86 (dd, J=7, 7 Hz, 2 H). MS calculated for C<sub>11</sub>H<sub>12</sub>F<sub>3</sub>NO<sub>2</sub>+H: 248, observed: 248.

*N-Trifluoroacetyl-2-iodo-5-methoxyphenethylamine*

A solution of N-trifluoroacetyl-3-methoxyphenethylamine (15.8 g, 64 mmol) in methanol (325 mL) was cooled to -78 C, and treated with CaCO<sub>3</sub> (14.7 g, 145 mmol), followed by a solution of ICl (29 g, 181 mmol) in methanol (40 mL). The reaction was allowed to warm to 20 C while stirring overnight and then filtered, concentrated, dissolved in EtOAc (200 mL), washed twice with 5% aqueous sodium bisulfite (100 mL), once with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 23.8 g of a white solid powder. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.68 (d, J=9 Hz, 1 H), 6.76 (s, 1 H), 6.57 (d, J=9 Hz, 1 H), 6.42 (bs, 1 H), 3.77 (s, 3 H), 3.61 (dd, J=7, 7 Hz, 2 H), 2.99 (dd, J=7, 7 Hz, 2 H). MS calculated for C<sub>11</sub>H<sub>11</sub>F<sub>3</sub>INO<sub>2</sub>+H: 374, observed: 374.

*N-Allyl, N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine*

A solution of N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (23.8 g, 63.8 mmol) in toluene (425 mL) was sequentially treated with K<sub>2</sub>CO<sub>3</sub> (12.4 g, 89.8 mmol), KOH (11.6 g, 207 mmol), n-Bu<sub>4</sub>NBr (2.2 g, 6.9 mmol) and allyl bromide (10.7 g, 89.8 mmol). The mixture was stirred at 80 C for 3.5 hours, cooled to 20 C, acidified with 10% aqueous HCl, separated and the aqueous phase extracted with ether (500 mL). The combined organic phases were washed with brine (200 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 20.5 g of a brown oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), mixture of rotamers d 7.67 (m, 1 H), 6.80 (m, 1 H), 6.57 (m, 1 H), 5.9-5.6 (bm, 1 H), 5.27 (m, 2 H), 4.11 (d, J=6 Hz, 0.5 H), 3.85 (d, J=6 Hz, 0.5 H), 3.77 (m, 3 H), 3.55 (m, 2 H), 3.00 (m, 2 H). MS calculated for C<sub>14</sub>H<sub>15</sub>F<sub>3</sub>INO<sub>2</sub>+H: 414, observed: 414.

*N-Trifluoroacetyl-7-methoxy-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-allyl, N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (20.5 g, 50 mmol) in dimethylformamide (250 mL) is treated with KOAc (14.6 g, 149 mmol), n-Bu<sub>4</sub>NBr (16.0 g, 50 mmol), PPh<sub>3</sub> (1.3 g, 5.0 mmol), Pd(OAc)<sub>2</sub> (0.56 g, 2.5 mmol) and stirred overnight at 90 C. The product mixture was cooled to 20 C, filtered, diluted with water (500 mL) and extracted with ether (3 x 500 mL). The combined organic phases were washed with water (100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 6.6 g of a yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) d 7.26 (d, J=8 Hz, 1 H), 6.77 (d, J=8 Hz, 1 H), 6.66 (s, 1 H), 5.34-5.19 (m, 2 H), 4.40 (m, 2 H), 3.83 (m, 2 H), 3.80 (s, 3 H), 3.00 (m, 2 H). MS calculated for C<sub>14</sub>H<sub>14</sub>F<sub>3</sub>NO<sub>2</sub>+H: 285, observed: 285.

*N-Trifluoroacetyl-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-methoxy-1-methylene-2,3,4,5-trihydro-1H-3-benzazepine (6.6 g, 23.2 mmol) in ethanol (100 mL), was treated with 10% Pd/C (0.75 g, 2.3 mmol) and stirred overnight under an atmosphere of hydrogen.

The product mixture was filtered through a pad of celite and silica and the solvent removed to give 6.27 g of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.10 (m, 1 H), 6.74 (m, 1 H), 6.68 (m, 1 H), 4.1-3.8 (bm, 2 H), 3.8 (s, 3H), 3.5 (m, 1.5 H), 3.4 (m, 0.5 H), 3.2-2.9 (bm, 4 H), 1.32 (m, 3 H). MS calculated for C<sub>14</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>2</sub>+H: 288, observed: 288.


*N-Trifluoroacetyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (1.25 g, 4.35 mmol) in acetonitrile (40 mL) was treated with N-bromosuccinimide (0.852 g, 4.79 mmol) and stirred overnight at 20 C. The product mixture was diluted with EtOAc (200 mL), washed with saturated aqueous sodium bisulfite (100 mL) and brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 1.55 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.34 (s, 1 H), 6.65 (m, 1 H), 3.87 (s, 3 H), 3.81 (m, 1 H), 3.55 (m, 1.3 H), 3.37 (m, 0.7 H), 3.2-2.9 (bm, 4 H), 1.30 (m, 3 H). MS calculated for C<sub>14</sub>H<sub>15</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 366, observed: 366.

*8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.95 g, 2.59 mmol) in methanol (20 mL) was treated with 15% aqueous NaOH (25 mL), and stirred overnight at 20 C. The product mixture was diluted with water (100 mL), extracted twice with EtOAc (100 mL), the combined organic phases were washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.687 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.92 (s, 1 H), 6.34 (s, 1 H), 3.87 (s, 3 H), 3.1-2.9 (m, 6 H), 2.75 (m, 1 H), 2.60 (bs, 1 H), 1.31 (d, J=7 Hz, 3 H). MS calculated for C<sub>12</sub>H<sub>16</sub>BrNO+H: 270, observed: 270.

**Example 2: (R,S) 8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-8-chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of *N*-trifluoroacetyl-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.900 g, 2.67 mmol) in acetonitrile (30 mL) was treated with *N*-chlorosuccinimide (0.357 g, 2.67 mmol) and stirred overnight at 70 C. The product mixture was diluted with water (100 mL), extracted twice with EtOAc (100 mL), the combined organic phases washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (20% EtOAc in hexane, silica) resulted in 0.399 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.17 (s, 1 H), 6.68 (m, 1 H), 3.88 (s, 3 H), 3.78 (m, 1 H), 3.6-3.3 (m, 2 H), 3.2-2.9 (m, 4 H), 1.34 (m, 3 H). MS calculated for C<sub>14</sub>H<sub>15</sub>ClF<sub>3</sub>NO<sub>2</sub>+H: 322, observed: 322.

*8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of *N*-trifluoroacetyl-8-chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.399 g, 1.24 mmol) in methanol (20 mL) was treated with 15% aqueous NaOH (20 mL), and stirred overnight at 20 C. The product mixture was diluted with water (100 mL), extracted twice with EtOAc (100 mL), the combined organic phases were washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.306 g of a yellow solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.05 (s, 1 H), 6.59 (s, 1 H), 3.80 (s, 3 H), 3.0-2.8 (m, 6 H), 2.62 (m, 1 H), 2.16 (bs, 1 H), 1.24 (d, J=7 Hz, 3 H). MS calculated for C<sub>12</sub>H<sub>16</sub>ClNO+H: 226, observed: 226.

**Example 3: (R,S) 8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (1.50 g, 5.22 mmol) in methanol (70 mL) was treated with  $\text{CaCO}_3$  (1.06 g, 10.44 mmol) followed by a solution of  $\text{ICl}$  (1.70 g, 10.44 mmol) in methanol (10 mL), and stirred overnight at 20 C. The product mixture was filtered, concentrated, dissolved in EtOAc (200 mL), extracted twice with 5% aqueous sodium bisulfite (100 mL), once with brine (100 mL), dried with  $\text{Na}_2\text{SO}_4$  and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 1.54 g of a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , mixture of rotamers)  $\delta$  7.55 (m, 1 H), 6.57 (m, 1 H), 3.86 (s, 3 H), 3.80 (m, 1 H), 3.60-3.30 (m, 2 H), 3.20-2.80 (m, 4 H), 1.30 (m, 3 H). MS calculated for  $\text{C}_{14}\text{H}_{15}\text{F}_3\text{INO}_2+\text{H}$ : 414, observed: 414.

*8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.600 g, 1.45 mmol) in methanol (20 mL) was treated with 15% aqueous NaOH (20 mL), and stirred for 3 hours at 50 C. The product mixture was diluted with water (100 mL), extracted twice with EtOAc (100 mL), the combined organic phases were washed with brine (100 mL), dried with  $\text{Na}_2\text{SO}_4$  and concentrated to give 0.425 g of a yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.52 (s, 1 H), 6.57 (s, 1 H), 3.86 (s, 3 H), 3.12-3.06 (m, 4 H), 2.95 (m, 2 H), 2.75 (m, 1 H), 2.43 (bs, 1 H), 1.33 (d,  $J=8$  Hz, 3 H). MS calculated for  $\text{C}_{12}\text{H}_{16}\text{INO}+\text{H}$ : 318, observed: 318.

**Example 4: (R,S) 8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (1.50 g, 4.10 mmol) in dichloromethane (80 mL) was treated dropwise with BBr<sub>3</sub> (9.4 mL of a 1.0M solution in CH<sub>2</sub>Cl<sub>2</sub>, 9.4 mmol), and the mixture stirred overnight while warming to 20 C. The excess BBr<sub>3</sub> was quenched with the dropwise addition of water, the mixture diluted with ether (200 mL), washed with Na<sub>2</sub>CO<sub>3</sub> (100 mL) and brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 1.25 g of a white solid foam. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.25 (s, 1 H), 6.79 (m, 1 H), 3.79 (m, 1 H), 3.7-3.3 (m, 2 H), 3.2-2.8 (m, 4 H), 1.32 (m, 3 H).

*8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.655 g, 1.89 mmol) in methanol (20 mL) was treated with 15% aqueous NaOH (20 mL), and stirred overnight at 20 C. The product mixture was diluted with water (100 mL), extracted twice with EtOAc (100 mL), the combined organic phases were washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.460 g of a clear oil. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.11 (s, 1 H), 6.65 (s, 1 H), 2.90 (m, 1 H), 2.73 (m, 5 H), 2.55 (m, 1 H), 1.19 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>14</sub>BrNO+H: 256, observed: 256.

**Example 5: (R,S) 7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-7-allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.150 g, 0.426 mmol) in dichloromethane (5 mL) was treated with allyl bromide (0.155 g, 1.28 mmol) and DBU (0.195 g, 1.28 mmol) and then stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (50 mL), washed with 5% aqueous HCl (20 mL), brine (20 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.149 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.34 (s, 1 H), 6.65 (m, 1 H), 6.04 (m, 1 H), 5.47 (d, J=17 Hz, 1 H), 5.30 (d, J=9 Hz, 1 H), 4.59 (s, 2 H), 3.80 (m, 1 H), 3.6-3.3 (m, 3 H), 3.2-2.8 (m, 4 H), 1.31 (m, 3 H). MS calculated for C<sub>16</sub>H<sub>17</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 392, observed: 392.

*7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (1.18 g, 3.00 mmol) in methanol (35 mL) was treated with 15% aqueous NaOH (35 mL), and stirred overnight at 20 C. The product mixture was diluted with water (200 mL), extracted twice with EtOAc (200 mL), the combined organic phases were washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.880 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (s, 1 H), 6.63 (s, 1 H), 6.04 (m, 1 H), 5.47 (d, J=17 Hz, 1 H), 5.29 (d, J=11 Hz, 1 H), 4.58 (s, 2 H), 3.01 (m, 3 H), 2.89 (m, 3 H), 2.75 (m, 1 H), 1.31 (d, J=7 Hz, 3 H). MS calculated for C<sub>14</sub>H<sub>18</sub>BrNO+H: 296, observed: 296.

**Example 6: (R,S) 7-Benzyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-7-benzyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine


A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.075 g, 0.213 mmol) in dichloromethane (5 mL) was treated with benzyl bromide (0.072 g, 0.64 mmol), DBU (0.100 g, 0.64 mmol), and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (50 mL), washed with 5% aqueous HCl (20 mL), brine (20 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.081 g of a clear oil. MS calculated for C<sub>20</sub>H<sub>19</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 442, observed: 442.

7-Benzyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-7-benzyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.81 g, 1.83 mmol) in methanol (20 mL) was treated with 15% aqueous NaOH (20 mL), and stirred overnight at 20 C. The product mixture was diluted with water (200 mL), extracted twice with EtOAc (200 mL), the combined organic phases were washed with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.412 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.38 (d, J=8 Hz, 2 H), 7.30 (dd, J=7, 8 Hz, 2 H), 7.23 (m, 2 H), 6.61 (s, 1 H), 5.03 (s, 2 H), 2.94 (m, 3 H), 2.81 (m, 3 H), 2.62 (m, 1 H), 2.30 (bs, 1 H), 1.24 (d, J=7 Hz, 3 H). MS calculated for C<sub>18</sub>H<sub>20</sub>BrNO+H: 346, observed: 346.



**Example 7: (R,S) 8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-8-bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.015 g, 0.043 mmol) in dichloromethane (1 mL) was treated with ethyl iodide (0.016 g, 0.102 mmol), DBU (0.016 g, 0.102 mmol) and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (10 mL), washed with 5% aqueous HCl (5 mL), brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.010 g of a clear oil.

*8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.010 g, 0.026 mmol) in methanol (1 mL) was treated with 15% aqueous NaOH (1 mL), and stirred overnight at 20 C. The product mixture was diluted with water (3 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (3 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.007 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (s, 1 H), 6.63 (s, 1 H), 4.07 (q, J=6 Hz, 2 H), 3.03 (m, 3 H), 2.91 (m, 3 H), 2.73 (m, 1 H), 2.26 (bs, 1 H), 1.46 (t, J=6 Hz, 3 H), 1.32 (d, J=7 Hz, 3 H). MS calculated for C<sub>15</sub>H<sub>17</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 380, observed: 380.

**Example 8: (R,S) 8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N-Trifluoroacetyl-8-bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.035 g, 0.099 mmol) in dichloromethane (1 mL) was treated with isopropyl bromide (0.037 g, 0.297 mmol), DBU (0.048 g, 0.205 mmol) and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (10 mL), washed with 5% aqueous HCl (5 mL), brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.014 g of a clear oil. MS calculated for C<sub>16</sub>H<sub>19</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 394, observed: 394.

*8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.014 g, 0.035 mmol) in methanol (1 mL) was treated with 15% aqueous NaOH (1 mL), and stirred overnight at 20 C. The product mixture was diluted with water (3 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (3 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.008 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 (s, 1 H), 6.64 (s, 1 H), 4.48 (m, 1 H), 2.98 (m, 3 H), 2.87 (m, 3 H), 1.36 (m, 6 H), 1.30 (d, J=7 Hz, 3 H). MS calculated for C<sub>14</sub>H<sub>20</sub>BrNO+H: 298, observed: 298.


**Example 9: (R,S) N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



A solution of 8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (6 mg, .022 mmol) in methanol (1 mL) was treated with excess paraformaldehyde, 1.0 M HCl in ether (0.004 mL, 0.004 mmol), NaBH<sub>3</sub>CN (1.0 mg, 0.013 mmol), and stirred overnight at 20 C. The product mixture was diluted


with 5% aqueous NaOH (5 mL), extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL each), the combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, silica) resulted in 5 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.31 (s, 1 H), 6.66 (s, 1 H), 3.87 (s, 3 H), 3.26 (bm, 2 H), 3.01 (bs, 1 H), 2.85 (m, 2 H), 2.45 (s, 3 H), 2.45-2.25 (m, 2 H), 1.36 (d, J=7 Hz, 3 H). MS calculated for C<sub>13</sub>H<sub>18</sub>BrNO+H: 284, observed: 284.

**Example 10: (R,S) N-Propyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



A solution of 8-Bromo-7-methoxy-1-methyl-1,2,4,5-tetrahydro-3H-3-benzazepine (6 mg, .022 mmol) in methanol (1 mL) was treated with propionaldehyde (5.0 mg, 0.067 mmol), 1.0 M HCl in ether (0.004 mL, 0.004 mmol), NaBH<sub>3</sub>CN (1.0 mg, 0.013 mmol), and stirred overnight at 20 C. The product mixture was diluted with 5% aqueous NaOH (5 mL), extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL each), the combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, silica) resulted in 4 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.33 (s, 1 H), 6.87 (s, 1 H), 3.84 (s, 3 H), 3.25 (m, 2 H), 3.11 (m, 2 H), 2.97 (m, 1 H), 2.78 (bm, 2 H), 2.63 (bm, 2 H), 1.67 (m, 2 H), 1.38 (d, J=7 Hz, 3 H), 0.96 (t, J=7 Hz, 3 H). MS calculated for C<sub>15</sub>H<sub>22</sub>BrNO+H: 312, observed: 312.

**Example 11: (R,S) 7-Hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




*N*-Trifluoroacetyl-7-hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (80 mg, 0.19 mmol) in dichloromethane (3 mL) was treated with BBr<sub>3</sub> (0.40 mL of a 1.0M solution in CH<sub>2</sub>Cl<sub>2</sub>, 0.40 mmol) and stirred overnight at 20 C. The excess BBr<sub>3</sub> was quenched with water and the product mixture was diluted with ether (20 mL), washed with Na<sub>2</sub>CO<sub>3</sub> (10 mL) and brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 74 mg of a white solid. MS calculated for C<sub>13</sub>H<sub>13</sub>F<sub>3</sub>INO<sub>2</sub>+H: 400, observed: 400.

*7-Hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (25 mg, 0.063 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 13 mg of a white solid. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.46 (s, 1 H), 6.64 (s, 1 H), 3.16 (m, 3 H), 2.94 (m, 3 H), 2.81 (m, 1 H), 1.35 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>14</sub>INO+H: 304, observed: 304.

**Example 12: (R,S) 7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine**



*N-Trifluoroacetyl-7-allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*


A solution of N-trifluoroacetyl-7-hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine (30 mg, 0.075 mmol) in dichloromethane (2 mL) was treated with allyl bromide (18 mg, 0.15 mmol), DBU (23 mg, 0.15 mmol) and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (10 mL), washed with 5% aqueous HCl (5 mL), brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and

concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 23 mg of a clear oil. MS calculated for  $C_{16}H_{17}F_3INO_2+H$ : 440, observed: 440.

*7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

5 A solution of N-trifluoroacetyl-7-allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (23 mg, 0.058 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with  
10  $Na_2SO_4$  and concentrated to give 18 mg of a white solid. MS calculated for  $C_{14}H_{18}INO+H$ : 344, observed: 344.

**Example 13: (R,S) 3,5-Dimethyl-6,7,8,9-tetrahydro-5H-1-oxa-7-aza-cycloheptaindene**



*N-trifluoroacetyl-3,5-dimethyl-6,7,8,9-tetrahydro-5H-1-oxa-7-aza-cycloheptaindene*


15 A solution of N-trifluoroacetyl-7-allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (158 mg, 0.360 mmol) in dimethylformamide (4 mL) was treated with KOAc (106 mg, 1.08 mmol), *n*-Bu<sub>4</sub>NBr (116 mg, 0.360 mmol), PPh<sub>3</sub> (13 mg, 0.036 mmol), Pd(OAc)<sub>2</sub> (4 mg, 0.018 mmol) and stirred  
20 overnight at 100 C. The product mixture was filtered, water (10 mL) added and then extracted twice with EtOAc (10 mL). The combined organic phases were washed with brine (10 mL), dried with  $Na_2SO_4$  and concentrated. Flash  
25 chromatography (5% EtOAc in hexane, silica) resulted in 15 mg of a clear oil. MS calculated for  $C_{16}H_{16}F_3NO_2+H$ : 312, observed: 312.

*3,5-Dimethyl-6,7,8,9-tetrahydro-5H-1-oxa-7-aza-cycloheptaindene*

A solution of N-trifluoroacetyl-3,5-dimethyl-6,7,8,9-tetrahydro-5H-1-oxa-

7-aza-cycloheptaindene (15 mg, 0.048 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 10 mg of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.25 (s, 1 H), 7.12 (s, 1 H), 7.09 (s, 1 H), 3.12 (m, 1 H), 2.97 (m, 4 H), 2.85 (m, 1 H), 2.64 (bm, 1 H), 2.15 (s, 3 H), 1.34 (d, J=8 Hz, 3 H). MS calculated for C<sub>14</sub>H<sub>17</sub>NO+H: 216, observed: 216.

**Example 14: (R,S) 7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-8-chloro-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of *N*-trifluoroacetyl-8-chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (48 mg, 0.15 mmol) in dichloromethane (2 mL) was treated with BBr<sub>3</sub> (0.30 mL of a 1.0M solution in CH<sub>2</sub>Cl<sub>2</sub>, 0.30 mmol) and stirred overnight at 20 C. The excess BBr<sub>3</sub> was quenched with water and the resulting mixture diluted with ether (20 mL), washed with Na<sub>2</sub>CO<sub>3</sub> (10 mL) and brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 24 mg of a white solid. MS calculated for C<sub>13</sub>H<sub>13</sub>ClF<sub>3</sub>NO<sub>2</sub>+H: 308, observed: 308.

*N*-Trifluoroacetyl-7-allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of *N*-trifluoroacetyl-8-chloro-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (24 mg, 0.078 mmol) in dichloromethane (2 mL) was treated with allyl bromide (18 mg, 0.15 mmol), DBU (23 mg, 0.15 mmol) and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (10 mL),


washed with 5% aqueous HCl (5 mL), brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 23 mg of a white solid. MS calculated for C<sub>16</sub>H<sub>17</sub>ClF<sub>3</sub>NO<sub>2</sub>+H: 348, observed: 348.

5 *7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (23 mg, 0.066 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5  
10 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 19 mg of a white solid. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.12 (s, 1 H), 6.81 (s, 1 H), 6.03 (m, 1 H), 5.43 (d, J=17 Hz, 1 H), 5.24 (d, J=10 Hz, 1 H), 4.57 (d, J=5 Hz, 2 H), 3.1-2.9 (m, 5 H), 2.81 (m, 1 H), 2.63 (m, 1 H), 1.30 (d, J=7 Hz, 3 H).

15

**Example 15: (R,S) 7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1H-3-benzazepine**




20 *N-Trifluoroacetyl-7-methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoromethylacetyl-8-bromo-7-methoxy-1-methyl-1,2,4,5-tetrahydro-3H-3-benzazepine (51 mg, 0.14 mmol) in 1,4-dioxane (2 mL) was treated with thiophene-2-boronic acid (36 mg, 0.28 mmol), K<sub>2</sub>CO<sub>3</sub> (58 mg, 0.42 mmol), water (0.1 mL), Pd(PPh<sub>3</sub>)<sub>4</sub> (16 mg, 0.014 mmol) and stirred overnight  
25 at 100 C. The product mixture was diluted with EtOAc, filtered, absorbed on silica and purified by flash chromatography (10% EtOAc in hexane, silica) resulting in 28 mg of a yellow solid. MS calculated for C<sub>18</sub>H<sub>18</sub>F<sub>3</sub>NO<sub>2</sub>S+H: 370, observed: 370.

*7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (28 mg, 0.076 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred 0.5 hours at 50 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 18 mg of a yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.45 (d, J=4 Hz, 1 H), 7.39 (s, 1 H), 7.27 (d, J=6 Hz, 1 H), 7.07 (dd, J=4, 6 Hz, 1 H), 6.71 (s, 1 H), 3.90 (s, 3 H), 3.1-2.9 (m, 6 H), 2.80 (m, 1 H), 2.22 (bs, 1 H), 1.38 (d, J=7 Hz, 3 H). MS calculated for C<sub>16</sub>H<sub>19</sub>NOS+H: 274, observed: 274.

**Example 16: (R,S) 8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine***N-Trifluoroacetyl-8-cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (18 mg, 0.05 mmol) in dimethylformamide (1 mL) was treated with CuCN (20 mg, 0.24 mmol) and the mixture was microwaved at 200 C for 0.5 hours. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (35% EtOAc in hexane, silica) resulted in 10 mg of a clear oil. MS calculated for C<sub>15</sub>H<sub>15</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>+H: 313, observed: 313.


*8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (10 mg, 0.032 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred 1 hour at 50 C. The product



mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 6.0 mg of a white solid. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.33 (s, 1 H), 6.93 (s, 1 H), 3.91 (s, 3 H), 3.18-2.97 (m, 5 H), 2.80 (m, 1 H), 2.60 (m, 1 H), 1.33 (d, J=8 Hz, 3 H). MS calculated for C<sub>13</sub>H<sub>16</sub>N<sub>2</sub>O+H: 217, observed: 217.

**Example 17: (R,S) 8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of diethyl zinc (1 mL, 1 M in hexanes) in dichloromethane (1 mL) at 0 °C was treated with trifluoroacetic acid in dichloromethane (0.5 mL) and the mixture stirred for 15 min. Diiodomethane (0.280 g, 1.0 mmol) in dichloromethane (0.5 mL) was then added and stirred for 15 minutes. *N*-Trifluoroacetyl-7-methoxy-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine (0.075 g, 0.26 mmol) in dichloromethane (1 mL) was added and the mixture stirred for 30 minutes at 0 °C and then for 2 hours at 20 °C. The product mixture was quenched with aqueous saturated NH<sub>4</sub>Cl (5 mL), extracted twice with CH<sub>2</sub>Cl<sub>2</sub> (20 mL), washed with saturated aqueous NaHCO<sub>3</sub> (10 mL), washed with H<sub>2</sub>O (10 mL), and concentrated. Flash chromatography (7% EtOAc in hexanes, silica) resulted in 0.050 g of a white solid. MS calculated for C<sub>15</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>2</sub>+H: 300, observed: 300.

*N*-Trifluoroacetyl-8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine


A solution of *N*-trifluoroacetyl-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.025 g, 0.08 mmol) in acetonitrile (1 mL) was treated with *N*-bromosuccinimide (0.032 g, 0.18 mmol) and stirred for 2 hrs. at 50

C. The product mixture was concentrated and then purified by flash chromatography (10% EtOAc in hexanes, silica) resulting in 0.014 g of a white solid. MS calculated for  $C_{15}H_{15}BrF_3NO_2+H$ : 378, observed: 378.

5 *8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.014 g, 0.037 mmol) in methanol (1 mL) was treated with 15% aqueous NaOH (1 mL), and stirred for 2 hours at 50 C. The product mixture was diluted with brine (10 mL), extracted twice with EtOAc (10 mL), dried with  $MgSO_4$ , and concentrated to give 0.008 g of a clear oil. <sup>1</sup>H NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.26 (s, 1 H), 6.78 (s, 1 H), 3.83 (s, 3 H), 3.02 (m, 2 H), 2.92 (m, 2 H), 2.67 (s, 2 H), 0.91 (m, 2 H), 0.85 (m, 2 H). MS calculated for  $C_{13}H_{16}BrNO+H$ : 282, observed: 282.

15 **Example 18: (R,S) 8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N-Trifluoroacetyl-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

20 A solution of N-trifluoroacetyl-7-methoxy-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine (0.100 g, 0.35 mmol) in tetrahydrofuran (1 mL) was treated with  $BH_3$ -THF complex (0.36 mL, 1 M in THF), and stirred for 30 min. at 20 C. Water (0.5 mL), saturated aqueous  $NaHCO_3$  (0.5 mL), and 30%  $H_2O_2$  (0.2 mL) were added sequentially and the reaction stirred for 30 min. at 20 C. The product mixture was diluted with EtOAc (10 mL), washed with brine (10 mL), and concentrated. Flash chromatography (33% EtOAc in hexane, silica) resulted in 0.035 g of a clear oil. MS calculated for  $C_{14}H_{16}F_3NO_3+H$ : 304, observed: 304.


*N-Trifluoroacetyl-8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-*

*benzazepine*

A solution of N-trifluoroacetyl-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.035 g, 0.12 mmol) in acetonitrile (1 mL) was treated with N-bromosuccinimide (0.025 g, 0.14 mmol), and stirred for 30 min. at 20 C. The product mixture was concentrated and then purified by flash chromatography (33% EtOAc in hexane, silica) resulting in 0.019 g clear oil. MS calculated for  $C_{14}H_{15}BrF_3NO_3+H$ : 382, observed: 382.

*8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.009 g, 0.024 mmol) in methanol (1 mL) was treated with 15% aqueous NaOH (1 mL), and stirred for 1 hour at 50 C. The product mixture was diluted with brine (5 mL), extracted twice with EtOAc (5 mL), dried with  $MgSO_4$ , and concentrated to give 0.006 g clear oil.  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.28 (s, 1 H), 6.79 (s, 1 H), 3.84 (m, 2 H), 3.0-2.8 (m, 7 H). MS calculated for  $C_{12}H_{16}BrNO_2+H$ : 286, observed: 286.

**Example 19: (R,S) 8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine***N-Crotyl, N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine*

A solution of N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (6.68 g, 17.9 mmol) in toluene (100 mL) was treated with  $K_2CO_3$  (3.22 g, 23.3 mmol), KOH (3.01 g, 53.7 mmol),  $n-Bu_4NBr$  (0.580 g, 1.80 mmol) and crotyl bromide (3.15 g, 23.3 mmol). The mixture was stirred at 75 C for 16 hours, cooled to 20 C, diluted with  $Et_2O$  (500 mL), washed with 10% aqueous HCl (500 mL) and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 5.22 g of a clear oil. MS calculated for  $C_{15}H_{17}F_3INO_2+H$ : 428, observed: 428.

*N-Trifluoroacetyl-1-ethylene-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-crotyl, N-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (5.20 g, 12.2 mmol) in dimethylformamide (80 mL) was treated with KOAc (3.59 g, 36.6 mmol), n-Bu<sub>4</sub>NBr (3.93 g, 12.2 mmol), PPh<sub>3</sub> (0.320g, 1.22 mmol), Pd(OAc)<sub>2</sub> (0.137 g, 0.61 mmol) and stirred overnight at 90 C. The product mixture was cooled to 20 C, diluted with water (200 mL), extracted twice with ether (500 mL), the combined organic phases washed twice with brine (200 mL), and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 2.29 g of a clear oil, which consists of a mixture of olefinic isomers. MS calculated for C<sub>15</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>2</sub>+H: 300, observed: 300.

*N-Trifluoroacetyl-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-1-ethylene-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (2.29 g, 7.65 mmol) in methanol (100 mL) was treated with 10% Pd/C (4.0 g, 0.77 mmol) and stirred overnight under an atmosphere of hydrogen. The product mixture was filtered through a pad of celite and silica, and the solvent removed to give 2.14 g of a clear oil. MS calculated for C<sub>15</sub>H<sub>18</sub>F<sub>3</sub>NO<sub>2</sub>+H: 302, observed: 302.

*N-Trifluoroacetyl-8-bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*


A solution of N-trifluoroacetyl-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.710 g, 2.36 mmol) in acetonitrile (20 mL) was treated with N-bromosuccinimide (0.504 g, 2.83 mmol), and stirred overnight at 20 C. The product mixture was concentrated, diluted with EtOAc (100 mL), washed with water (50 mL) and brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc in hexanes, silica) resulted in 0.561 g of a clear oil. MS calculated for C<sub>15</sub>H<sub>17</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 380, observed: 380.

*8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1-ethyl-7-methoxy-2,3,4,5-

tetrahydro-1*H*-3-benzazepine (0.561 g, 1.48 mmol) in methanol (30 mL) was  
 treated with 15% aqueous NaOH (30 mL), and stirred overnight at 20 C. The  
 product mixture was diluted with brine (100 mL), extracted twice with EtOAc  
 (200 mL), dried with MgSO<sub>4</sub>, and concentrated to give 0.412 g of a clear oil. <sup>1</sup>H  
 NMR (400 MHz, CD<sub>3</sub>OD) δ 7.24 (s, 1 H), 6.76 (s, 1 H), 3.83 (s, 3 H), 3.02 (m, 3  
 H), 2.91 (s, 1 H), 2.85-2.76 (m, 3 H), 2.63 (m, 1 H), 1.78 (m, 1 H), 1.72 (m, 1 H),  
 0.94 (dd, J=8, 8 Hz, 3 H). MS calculated for C<sub>13</sub>H<sub>18</sub>BrNO+H: 284, observed: 284.

**Example 20: (R,S) 8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine**



*N*-Trifluoroacetyl-8-chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine


A solution of *N*-trifluoroacetyl-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-  
 3-benzazepine (0.600 g, 1.99 mmol) in acetonitrile (20 mL) was treated with *N*-  
 chlorosuccinimide (0.057 g, 0.32 mmol), and stirred overnight at 60 C. The  
 product mixture was concentrated, diluted with EtOAc (100 mL), washed with  
 water (50 mL) and brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash  
 chromatography (10% EtOAc in hexanes, silica) resulted in 0.421 g of a clear oil.  
 MS calculated for C<sub>15</sub>H<sub>17</sub>ClF<sub>3</sub>NO<sub>2</sub>+H: 336, observed: 336.

*8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of *N*-trifluoroacetyl-8-chloro-1-ethyl-7-methoxy-2,3,4,5-  
 tetrahydro-1*H*-3-benzazepine (0.421 g, 1.25 mmol) in methanol (30 mL) was  
 treated with 15% aqueous NaOH (30 mL), and stirred overnight at 20 C. The  
 product mixture was diluted with brine (100 mL), extracted twice with EtOAc  
 (200 mL), dried with MgSO<sub>4</sub>, and concentrated to give 0.241 g of a clear oil. <sup>1</sup>H  
 NMR (400 MHz, CD<sub>3</sub>OD) δ 7.05 (s, 1 H), 6.79 (s, 1 H), 3.84 (s, 3 H), 3.03 (m, 3  
 H), 2.91 (s, 1 H), 2.86-2.76 (m, 3 H), 2.64 (m, 1 H), 1.81 (m, 1 H), 1.72 (m, 1 H),

0.93 (dd, J=8, 8 Hz, 3 H). MS calculated for  $C_{13}H_{18}ClNO+H$ : 240, observed: 240.

**Example 21: (R,S) 8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-(3-methylbut-2-enyl), *N*-trifluoroacetyl-2-iodo-5-methoxyphenethylamine

A solution of *N*-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (0.700 g, 1.88 mmol) in toluene (25 mL) was treated with  $K_2CO_3$  (0.340 g, 2.4 mmol), KOH (0.210 g, 3.76 mmol), *n*-Bu<sub>4</sub>NBr (0.060 g, 0.19 mmol) and 4-bromo-2-methyl-2-butene (0.364 g, 2.44 mmol). The mixture was stirred at 80 C for 3 hours, cooled to 20 C, diluted with ether (100 mL), washed with 10% HCl (50 mL) and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 0.272 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.65 (m, 1 H), 6.75 (m, 1 H), 6.54 (m, 1 H), 5.20 (m, .4 H), 5.0 (m, .6 H), 4.10 (m, 1 H), 3.82 (m, 1 H), 3.76 (d, 2H), 3.50 (m, 2 H), 3.02 (m, 2H), 1.75 (m, 3H), 1.66(m, 3H).

*N*-Trifluoroacetyl-1-isopropylene-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of *N*-(3-methylbut-2-enyl), *N*-trifluoroacetyl-2-iodo-5-methoxyphenethylamine (0.272 g, 0.62 mmol) in dimethylformamide (12 mL) was treated with KOAc (0.183 g, 1.86 mmol), *n*-Bu<sub>4</sub>NBr (0.200 g, .062 mmol), PPh<sub>3</sub> (0.016g, 0.062 mmol), Pd(OAc)<sub>2</sub> (0.183 g, 1.86 mmol) and stirred overnight at 90 C. The product mixture was cooled to 20 C, diluted with water (50 mL), extracted twice with ether (50 mL), the combined organic phases were washed with brine (50 mL), and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 0.096 g of a clear oil. MS calculated for  $C_{16}H_{18}F_3NO_2+H$ : 314, observed: 314.

*N*-Trifluoroacetyl-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-1-isopropylene-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.096 g, 0.31 mmol) in ethanol (2 mL) was treated with 10% Pd/C (0.033 g, .031 mmol) and stirred overnight under an atmosphere of hydrogen. The product mixture was filtered through a pad of celite and silica, and the solvent removed to give 0.091 g of a clear oil. MS calculated for  $C_{16}H_{20}F_3NO_2+H$ : 316, observed: 316.


*N-Trifluoroacetyl-8-bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.091 g, 0.29 mmol) in acetonitrile (3 mL) was treated with N-bromosuccinimide (0.057 g, 0.32 mmol), and stirred overnight at 20 C. After removing the solvent, flash chromatography (10% EtOAc in hexanes, silica) resulted in 0.056 g of a clear oil. MS calculated for  $C_{16}H_{19}BrF_3NO_2+H$ : 394, observed: 394.

*8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine (0.013 g, 0.03 mmol) in methanol (0.5 mL) was treated with 15% aqueous NaOH (0.5 mL), and stirred overnight at 20 C. The product mixture was diluted with brine (5 mL), extracted twice with EtOAc (5 mL), dried with  $MgSO_4$ , and concentrated to give 0.10 g of a clear oil.  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.08 (s, 1 H), 6.64 (s, 1 H), 3.72 (s, 3 H), 3.2-3.10 (m, 3 H), 2.7-2.5 (m, 3 H), 2.3-2.1 (m, 2 H), 0.96 (d, 3 H), 0.63 (d, 3 H). MS calculated for  $C_{14}H_{20}BrNO+H$ : 298, observed: 298.

**Example 22: (R,S) 8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine**




*N-Trifluoroacetyl-8-bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.041 g, 0.10 mmol) in dichloromethane (1 mL) was treated with BBr<sub>3</sub> (0.32 ml, 1.0 M solution in CH<sub>2</sub>Cl<sub>2</sub>) and stirred overnight at 20 C. The excess BBr<sub>3</sub> is quenched with water and the resulting mixture diluted with ether (50 mL), washed twice with saturated aqueous Na<sub>2</sub>CO<sub>3</sub> (20 mL) and concentrated. Flash chromatography (20% EtOAc in hexanes, silica) resulted in 0.037 g clear oil. MS calculated for C<sub>15</sub>H<sub>17</sub>BrF<sub>3</sub>NO<sub>2</sub>+H: 380, observed: 380.

*8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.018 g, 0.047 mmol) in methanol (1 mL) was treated with 15% aqueous NaOH (1 mL), and stirred for 3 hours at 50 C. The product mixture was brought to pH 7-8 with 10% aqueous HCl, extracted three times with EtOAc (50 mL), dried with MgSO<sub>4</sub>, and concentrated to give 0.013 g of a white solid. <sup>1</sup>H NMR (400MHz, CD<sub>3</sub>OD) δ 7.10 (s, 1 H), 6.60 (s, 1 H), 3.30 (m, 1H), 3.2-3.0 (m, 2 H), 2.78 (m, 1H), 2.7-2.5 (m, 2H), 2.3-2.1 (m, 2 H), 1.05 (d, 3 H), 0.73 (d, 3 H). MS calculated for C<sub>13</sub>H<sub>18</sub>BrNO+H: 284, observed: 284.

**Example 23: (R,S) 7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N-Trifluoroacetyl-7-allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.017 g, 0.045 mmol) in dichloromethane (1 mL) was treated with N'''-tert-butyl-N, N, N', N', N'', N''-hexamethylphosphorimidic triamide (0.016 g, 0.068 mmol), allyl bromide (0.011 g, 0.09 mmol) and stirred for



3 hours at 20 C. The product mixture was diluted with 10% aqueous HCl, extracted twice with dichloromethane (20 mL), and concentrated. Flash chromatography (10% EtOAc in hexanes, silica) resulted in 0.011 g of a clear oil. MS calculated for  $C_{18}H_{21}BrF_3NO_2+H$ : 420, observed: 420.

5


*7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.011 g, 0.026 mmol) in methanol (0.5 mL) was treated with 15% aqueous NaOH (0.5 mL), and stirred for 3 hours at 50 C. The product mixture was diluted with brine (5 mL), extracted twice with EtOAc (5 mL), dried with  $MgSO_4$ , and concentrated to give 0.010 g of a clear oil.  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.09 (s, 1 H), 6.62 (s, 1 H), 5.94 (m, 1 H), 5.32 (dd, 1 H), 5.12 (dd, 1 H), 4.46 (d, 2H), 3.19 (m, 1 H), 3.05 (m, 2 H), 2.66 (m, 1 H), 2.5 (bm, 2 H), 2.3-2.1 (m, 2 H), 0.95 (d, 3 H), 0.63 (d, 3 H). MS calculated for  $C_{16}H_{22}BrNO+H$ : 324, observed: 324.

10

15

**Example 24: 8-Bromo-7-methoxy-1,4-dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



20

*N-Trifluoroacetyl-1-(3-methoxyphenyl)-2-propylamine*

A solution of 1-(3-methoxyphenyl)-2-propylamine (3.59 g, 21.7 mmol) in dichloromethane (75 mL) at 0 C, was treated with pyridine (2.1 mL, 28.2 mmol), trifluoroacetic anhydride (5.9 g, 28.2 mmol), and then stirred for 3 hours while warming to 20 C. The product mixture was diluted with EtOAc (300 mL), washed sequentially with 10% aqueous HCl (100 mL), water (100 mL), brine (100 mL), dried with  $Na_2SO_4$  and concentrated. Flash chromatography (20% EtOAc in hexane, silica) resulted in 4.29 g of a yellow solid.  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.17 (dd,  $J=8, 8$  Hz, 1 H), 6.76 (m, 3 H), 4.19 (m, 1 H), 3.77 (s, 3 H), 2.78 (m, 2 H), 1.21 (d,  $J=7$  Hz, 2 H).

25

*N-Trifluoroacetyl-1-(2-iodo-5-methoxyphenyl)-2-propylamine*

A solution of N-trifluoroacetyl-1-(3-methoxyphenyl)-2-propylamine (4.29 g, 15.7 mmol) in methanol (100 mL) was cooled to -78 C and treated with CaCO<sub>3</sub> (3.17 g, 31.4 mmol), followed by a solution of ICl (6.37 g, 39.3 mmol) in methanol (50 mL). The reaction was allowed to warm to 20 C while stirring overnight. The product mixture was filtered, concentrated, dissolved in EtOAc (200 mL), washed twice with 5% aqueous sodium bisulfite (100 mL), once with brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 6.72 g of a white solid powder. MS calculated for C<sub>12</sub>H<sub>13</sub>F<sub>3</sub>INO<sub>2</sub>+H: 388, observed: 388.

*N-Allyl, N-trifluoroacetyl-1-(2-iodo-5-methoxyphenyl)-2-propylamine*

A solution of N-trifluoroacetyl-1-(2-iodo-5-methoxyphenyl)-2-propylamine (6.09 g, 15.7 mmol) in toluene (450 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (2.82 g, 20.4 mmol), KOH (2.45 g, 47.1 mmol), n-Bu<sub>4</sub>NBr (0.506 g, 1.57 mmol) and allyl bromide (2.47 g, 20.4 mmol), and stirred overnight at 80 C. The product mixture was acidified with 10% aqueous HCl, separated, the aqueous phase extracted with ether (500 mL), the combined organic phases were washed with brine (200 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 4.45 g of a brown oil.

*N-Trifluoroacetyl-7-methoxy-4-methyl-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-allyl, N-trifluoroacetyl-1-(2-iodo-5-methoxyphenyl)-2-propylamine (4.45 g, 10.8 mmol) in dimethylformamide (120 mL) was treated with KOAc (3.17 g, 32.3 mmol), n-Bu<sub>4</sub>NBr (3.47 g, 10.8 mmol), PPh<sub>3</sub> (0.283 g, 1.08 mmol), Pd(OAc)<sub>2</sub> (0.242 g, 1.08 mmol) and stirred overnight at 80 C. The product mixture was cooled to 20 C, filtered, diluted with water (200 mL), extracted with ether (3 x 200 mL), the combined organic phases washed with water (100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 1.39 g of a yellow oil.

*N-Trifluoroacetyl-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-methoxy-4-methyl-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine (1.39 g, 4.64 mmol) in ethanol (40 mL) was treated with 10% Pd/C (0.49 g, 0.46 mmol) and stirred overnight under an atmosphere of hydrogen. The product mixture was filtered through a pad of celite and silica and then concentrated. Flash chromatography (20% EtOAc in hexane, silica) resulted in 0.77 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.06 (m, 1 H), 6.71 (m, 1 H), 6.63 (m, 1 H), 4.38 (bm, 1 H), 3.8 (s, 3H), 3.6 (m, 1 H), 3.25 (m, 1 H), 3.18 (bm, 2 H), 2.72 (m, 1 H), 1.34 (m, 3 H) 1.22 (m, 3 H).

*N-Trifluoroacetyl-8-bromo-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*


A solution N-trifluoroacetyl-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.452 g, 1.50 mmol) in acetonitrile (20 mL) was treated with N-bromosuccinimide (0.294 g, 1.65 mmol) and stirred overnight at 20 C. The product mixture was diluted with EtOAc (100 mL), washed with sodium bisulfite (50 mL) and brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (20% EtOAc in hexane, silica) resulted in a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.32 (s, 1 H), 6.62 (m, 1 H), 4.37 (m, 1 H), 3.87 (s, 3 H), 3.81 (m, 1 H), 3.28-3.10 (m, 3 H), 2.73 (m, 1 H), 1.31 (m, 3 H), 1.25 (m, 3 H).

*8-Bromo-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-bromo-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (21 mg, 0.055 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (10 mL), the combined organic phases were washed with brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 11 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (s, 1 H), 6.64 (s, 1 H), 3.88 (s, 3 H), 3.02 (m, 2 H), 2.89 (dd, J=9, 14 Hz, 1 H), 2.80 (m, 1 H), 2.67 (d, J=14 Hz, 1 H), 2.53 (dd, J=10, 13, 1 H) 1.30

(d, J=7 Hz, 3 H), 1.19 (d, J=6 Hz, 3 H). MS calculated for C<sub>13</sub>H<sub>18</sub>BrNO+H: 284, observed: 284.

**Example 25: 7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-8-bromo-1,4-dimethyl-7-hydroxy-2,3,4,5-tetrahydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-bromo-1,4-dimethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.383 g, 1.01 mmol) in dichloromethane (30 mL) was treated with BBr<sub>3</sub> (2.35 mL of a 1.0M solution in CH<sub>2</sub>Cl<sub>2</sub>, 2.35 mmol) and stirred overnight while warming to 20 C. The excess BBr<sub>3</sub> is quenched with water, and the resulting mixture was diluted with ether (100 mL), washed with saturated aqueous Na<sub>2</sub>CO<sub>3</sub> (50 mL) and brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.302 g of a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.22 (m, 1 H), 6.77 (m, 1 H), 5.34 (s, 1 H), 4.35 (m, 1 H), 3.62 (m, 1 H), 3.24 (m, 1 H), 3.13 (m, 2 H), 2.69 (m, 1 H), 1.31 (m, 3 H), 1.22 (m, 3 H).

*N*-Trifluoroacetyl-7-allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepine


A solution N-trifluoroacetyl-8-bromo-1,4-dimethyl-7-hydroxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.030 g, 0.082 mmol) in dichloromethane (2 mL) was treated with allyl bromide (0.030 g, 0.246 mmol), DBU (0.037 g, 0.246 mmol) and stirred 2 hours at 20 C. The product mixture was diluted with EtOAc (10 mL), washed with 5% aqueous HCl (2 mL), brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.028 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 7.32 (s, 1 H), 6.62 (m, 1 H), 6.02 (m, 1 H), 5.45 (d, J=17 Hz, 1 H), 5.30 (d, J=11 Hz, 1

H), 4.58 (s, 2 H), 4.36 (m, 1 H), 3.62 (m, 1 H), 3.23 (m, 1 H), 3.11 (m, 1 H), 2.81 (d, J=10 Hz, 1 H), 2.70 (m, 1 H), 1.34 (m, 3 H), 1.21 (m, 3 H).

*7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

5 A solution of N-trifluoroacetyl-7-allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.028 g, 0.069 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred for 3 hours at 20 C. The product mixture was diluted with water (10 mL), extracted twice with EtOAc (10 mL), the combined organic phases were washed with brine (10 mL), dried with  
10 Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.020 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.30 (s, 1 H), 6.64 (s, 1 H), 6.06 (m, 1 H), 5.47 (d, J=17 Hz, 1 H), 5.30 (d, J=11 Hz, 1 H), 4.56 (s, 2 H), 3.03 (m, 2 H), 2.90 (dd, J=9, 14 Hz, 1 H), 2.80 (m, 1 H), 2.65 (d, J=14 Hz, 1 H), 2.55 (dd, J=10, 14 Hz, 1 H), 1.77 (bs, 1 H), 1.30 (d, J=7 Hz, 3 H), 1.20 (d, J=6 Hz, 3 H).

**Example 26: (R,S) 8-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N-Trifluoroacetyl-4-chlorophenethylamine*

20 A solution of 4-chlorophenethylamine (1.0 g, 6.4 mmol) in dichloromethane (20 mL) was cooled to 0 C, treated with pyridine (1.0 mL, 12.8 mmol), trifluoroacetic anhydride (1.6 g, 7.7 mmol) and then stirred for 1 hour while warming to 20 C. The product mixture was diluted with EtOAc (100 mL), washed sequentially with 10% aqueous HCl (50 mL), water (50 mL), brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 1.6 g of a white solid.

*N-Trifluoroacetyl-2-iodo-4-chlorophenethylamine*

25 A solution of N-trifluoroacetyl-4-chlorophenethylamine (1.6 g, 6.4 mmol) in dichloromethane (20 mL) was treated with bis(pyridine)iodonium(I)tetrafluoroborate (2.6 g, 7.0 mmol), CF<sub>3</sub>SO<sub>3</sub>H (2.1 g,

14.1 mmol) and stirred overnight at 20 C. The product mixture was concentrated, dissolved in EtOAc (100 mL), washed twice with 5% aqueous sodium bisulfite (50 mL), twice with saturated aqueous NaHCO<sub>3</sub>, (50 mL) once with brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.94 g of a clear oil. MS calculated for C<sub>10</sub>H<sub>8</sub>ClF<sub>3</sub>INO+H: 378, observed: 378.

*N-Allyl, N-trifluoroacetyl-2-iodo-4-chlorophenethylamine*

A solution of N-trifluoroacetyl-2-iodo-4-chlorophenethylamine (0.94 g, 2.4 mmol) in toluene (25 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (0.43 g, 3.12 mmol), KOH (0.40 g, 7.2 mmol), n-Bu<sub>4</sub>NBr (0.077 g, 0.24 mmol) and allyl bromide (0.43 g, 3.6 mmol) sequentially. The mixture was stirred at 80 C for 3.5 hours, cooled to 20 C and acidified with 10% aqueous HCl. The phases were separated, the aqueous phase extracted with ether (100 mL), the combined organic phases were washed with brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 0.76 g of a clear oil. MS calculated for C<sub>13</sub>H<sub>12</sub>ClF<sub>3</sub>INO+H: 418, observed: 418.

*N-Trifluoroacetyl-8-chloro-1-methylene-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-allyl, N-trifluoroacetyl-2-iodo-4-chlorophenethylamine (0.76 g, 1.8 mmol) in dimethylformamide (20 mL) was treated with KOAc (0.53 g, 5.4 mmol), n-Bu<sub>4</sub>NBr (0.58 g, 1.8 mmol), PPh<sub>3</sub> (0.047 g, 0.18 mmol), Pd(OAc)<sub>2</sub> (0.041 g, 0.18 mmol) and stirred overnight at 105 C. The product mixture was cooled to 20 C, filtered, diluted with water (100 mL), extracted with ether (3 x 100 mL), the combined organic phases washed with water (100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10% EtOAc in hexane, silica) resulted in 0.228 g of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (s, 1 H), 7.18 (m, 1 H), 7.04 (m, 1 H), 5.38 (m, 2 H), 5.40 (d, J=16 Hz, 2 H), 3.80 (m, 2 H), 3.00 (m, 2 H). MS calculated for C<sub>13</sub>H<sub>11</sub>ClF<sub>3</sub>NO+H: 290, observed: 290.

*N-Trifluoroacetyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*


A solution of N-trifluoroacetyl-8-chloro-1-methylene-2,3,4,5-tetrahydro-1H-

3-benzazepine (0.16 g, 0.55 mmol) in methanol (10 mL) was treated with 10% Pd/C (0.02 g) and stirred 30 minutes under an atmosphere of hydrogen. The product mixture was filtered, concentrated and purified by flash chromatography (5% EtOAc in hexane, silica) resulting in 0.057 g of a white solid. MS calculated for  $C_{13}H_{13}ClF_3NO+H$ : 292, observed: 292.

*8-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (65 mg, 0.22 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred for 3.5 hours at 60 C. The product mixture was concentrated, extracted 3 times with  $CH_2Cl_2$  (5 mL), dried with  $Na_2SO_4$  and concentrated to give 35 mg of a clear oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.11 (s, 1 H), 7.05 (d, J=8 Hz, 1 H), 6.98 (d, J=8 Hz, 1 H), 3.1-2.9 (m, 6 H), 2.71 (m, 1 H), 2.68 (bs, 1 H), 1.32 (d, J=8 Hz, 3 H). MS calculated for  $C_{11}H_{14}ClN+H$ : 196, observed: 196.

**Example 27: (R,S) 7-(2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N-Trifluoroacetyl -7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (0.506 g, 1.76 mmol) in dichloromethane (20 mL) was treated with  $BBr_3$  (4.1 mL of a 1.0M solution in  $CH_2Cl_2$ , 4.1 mmol) and stirred overnight while warming to 20 C. The excess  $BBr_3$  was quenched with water, and the resulting mixture was diluted with ether (200 mL), washed with  $Na_2CO_3$  (100 mL) and brine (100 mL), dried with  $Na_2SO_4$  and concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 0.460 g of a white solid foam. MS calculated for  $C_{13}H_{14}F_3NO_2+H$ : 274, observed: 274.

*N-Trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine,  
O-trifluoromethanesulfonate*

5 A solution of N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (460 mg, 1.76 mmol) in dichloromethane (15 mL) was treated with pyridine (417 mg, 5.27 mmol), trifluoromethanesulfonic anhydride (991 mg, 3.52 mmol) and stirred 1.5 hours at 20 C. The product mixture was diluted with dichloromethane (100 mL), washed with water (50 mL), 5% aqueous HCl (50 mL), saturated aqueous NaHCO<sub>3</sub> (50 mL), brine (50 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and  
10 concentrated. Flash chromatography (15% EtOAc in hexane, silica) resulted in 658 mg of a clear oil. MS calculated for C<sub>14</sub>H<sub>13</sub>F<sub>6</sub>NO<sub>4</sub>S+H: 406, observed: 406.

*N-Trifluoroacetyl-7-(2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

15 To a solution of N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, O-trifluoromethanesulfonate (100 mg, 0.25 mmol) in dimethylformamide (2 mL) was treated with (2-methyl-2H-pyrazol-3-yl)-tri-*n*-butyltin (138 mg, 0.37 mmol), LiCl (21 mg, 0.50 mmol), Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (35 mg, 0.05 mmol) and stirred at 100 C for 4 hours. The product mixture was diluted  
20 with EtOAc (20 mL), washed twice with water (10 mL), once with brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (30% EtOAc in hexane, silica) resulted in 80 mg of a clear oil. MS calculated for C<sub>17</sub>H<sub>18</sub>F<sub>3</sub>N<sub>3</sub>O+H: 338, observed: 338.

25 *7-(2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*


A solution of N-trifluoroacetyl-7-(2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (48 mg, 0.14 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and the solution stirred overnight at 20 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL),  
30 dried with Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated. Flash chromatography (0-15% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, silica) resulted in 30 mg of a clear oil. <sup>1</sup>H NMR (400 MHz,



CDCl<sub>3</sub>) d 7.48 (s, 1 H), 7.21 (m, 2 H), 7.13 (s, 1 H), 6.27 (s, 1 H), 3.89 (s, 3 H), 3.3-2.9 (m, 9 H), 2.79 (dd, J=7, 14 Hz, 1 H), 1.40 (d, J=8 Hz, 3 H). MS calculated for C<sub>15</sub>H<sub>19</sub>N<sub>3</sub>+H: 242, observed: 242.

5

**Example 28: (R,S) 7-(4-Bromo-2-methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-7-(4-bromo-2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

10

To a solution of *N*-trifluoroacetyl-7-(2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (30 mg, 0.082 mmol) in dichloromethane (1 mL) was treated with *N*-bromosuccinimide (15.3 mg, 0.086 mmol) and stirred overnight at 20 C. The product mixture was absorbed on silica and purified by flash chromatography (2-5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, silica) resulting in 37 mg of a white crystalline solid. MS calculated for C<sub>17</sub>H<sub>17</sub>BrF<sub>3</sub>N<sub>3</sub>O+H: 416, observed: 416.

15

*7-(4-Bromo-2-methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*


20

A solution of *N*-trifluoroacetyl-7-(4-bromo-2-Methyl-2H-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (37 mg, 0.089 mmol) in methanol (2 mL) was treated with of 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated. Flash chromatography (0-15% MeOH in CH<sub>2</sub>Cl<sub>2</sub>, silica) resulted in 28 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) d 7.50 (s, 1 H), 7.25 (d, J=8 Hz, 1 H), 7.17 (d, J=8 Hz, 1 H), 7.10 (s, 1 H), 3.83 (s, 3 H), 3.17 (m, 1 H), 3.1-2.9 (m, 8 H), 2.80 (dd, J=7, 13 Hz, 1 H), 2.48 (bs,

25


1 H), 1.40 (d, J=8 Hz, 3 H). MS calculated for  $C_{15}H_{18}BrN_3+H$ : 320, observed: 320.

5      **Example 29: (R,S) 7-(3-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




A solution of N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, *O*-trifluoromethanesulfonate (50 mg, 0.123 mmol) in 1,4-dioxane  
10 (1.5 mL) was treated with 2-chlorophenylboronic acid (39 mg, 0.243 mmol), CsF (56 mg, 0.37 mmol), water (50 mg, 2.78 mmol),  $Pd(PPh_3)_4$  (29 mg, 0.025 mmol) and stirred overnight at 75 C. The product mixture was diluted with EtOAc (20 mL), washed with water (10 mL), brine (10 mL), dried with  $Na_2SO_4$  and concentrated. Flash chromatography (10-20% EtOAc in hexane, silica) resulted in  
15 45 mg of a clear oil. MS calculated for  $C_{19}H_{17}ClF_3NO+H$ : 368, observed: 368. The product (27 mg, 0.073 mmol) was dissolved in methanol (2 mL) treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was concentrated, extracted 3 times with  $CH_2Cl_2$  (5 mL), dried with  $Na_2SO_4$  and the solvent evaporated to give 18 mg of a clear oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  
20 d 7.54 (s, 1 H), 7.42 (d, J=6 Hz, 1 H), 7.35-7.21 (m, 5 H), 3.14 (m, 1 H), 3.1-2.9 (m, 8 H), 2.80 (bm, 2 H), 1.38 (d, J=8 Hz, 3 H). MS calculated for  $C_{17}H_{18}ClN_3+H$ : 272, observed: 272.

**Example 30: (R,S) 7-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



A solution of N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, *O*-trifluoromethanesulfonate (50 mg, 0.123 mmol) in 1,4-dioxane (1.5 mL) was treated with 2-chlorophenylboronic acid (39 mg, 0.243 mmol), CsF (56 mg, 0.37 mmol), water (50 mg, 2.78 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (29 mg, 0.025 mmol) and stirred overnight at 75 C. The product mixture was diluted with EtOAc (20 mL), washed with water (10 mL), brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10-20% EtOAc in hexane, silica) resulted in 36 mg of a clear oil. MS calculated for C<sub>19</sub>H<sub>17</sub>ClF<sub>3</sub>NO+H: 368, observed: 368. The product (27 mg, 0.073 mmol) was dissolved in methanol (2 mL) treated with 15% aqueous NaOH (2 mL), and stirred overnight at 20 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated to give 24 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.44 (d, J=8 Hz, 1 H), 7.35-7.22 (m, 5 H), 7.15 (s, 1 H), 3.14 (m, 1 H), 3.1-2.9 (m, 8 H), 2.80 (dd, J=13, Hz, 1 H), 2.51 (bs, 1 H), 1.38 (d, J=8 Hz, 3 H). MS calculated for C<sub>17</sub>H<sub>18</sub>ClN<sub>3</sub>+H: 272, observed: 272.

**Example 31: (R,S) 8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-8-chloro-1-oxo-,3,4,5-trihydro-1H-3-benzazepine

A solution of N-trifluoroacetyl-8-chloro-1-methylene-3,4,5-trihydro-1H-3-benzazepine (0.23 g, 0.80 mmol) in 1:1 methanol/dichloromethane (45 mL) was cooled to -78 C, treated with ozone until the solution turned blue (about 20 minutes), PPh<sub>3</sub> (0.21 g, 0.80 mmol) was added and the resulting solution was

stirred 90 minutes while warming to 20 C. The product mixture was concentrated and purified by flash chromatography (30% EtOAc in hexane, silica) resulting in 0.215 g of a white solid. MS calculated for  $C_{12}H_9ClF_3NO_2 + H$ : 292, observed: 292.

5


*8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-chloro-1-oxo-3,4,5-trihydro-1H-3-benzazepine

(50 mg, 0.17 mmol) in methanol (2 mL) was treated with  $NaBH_4$  and the resulting mixture was stirred 16 hours at 20 C. The white solid product was collected by filtration, washed with water and dried, resulting in 30 mg of a white solid.  $^1H$  NMR (400 MHz,  $CD_3OD$ )  $\delta$  7.39 (s, 1 H), 7.12 (d,  $J=8$  Hz, 1 H), 7.06 (d,  $J=8$  Hz, 1 H), 4.74 (d,  $J=8$  Hz, 1 H), 3.1-2.7 (m, 6 H). MS calculated for  $C_{10}H_{12}ClNO + H$ : 198, observed: 198.

15


**Example 32: (R,S) 8-Bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



By the same general procedure as in example 26, (R,S) 8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 4-bromophenethylamine as a colorless oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.27 (s, 1 H), 7.22 (d,  $J=8$  Hz, 1 H), 6.94 (d,  $J=8$  Hz, 1 H), 3.1-2.85 (m, 6 H), 2.72 (m, 1 H), 2.25 (bs, 1 H), 1.33 (d,  $J=7$  Hz, 3 H). MS calculated for  $C_{11}H_{14}BrN + H$ : 240, observed: 240.

20

**Example 33: (R,S) 8-Fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




25

By the same general procedure as in example 26, (R,S) 8-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 4-fluorophenethylamine as a colorless oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.00 (dd,  $J=8, 10$  Hz, 1 H), 6.86


(d, J=10 Hz, 1 H), 6.76 (d, J=8 Hz, 1 H), 3.08-2.56 (m, 7 H), 1.85 (bs, 1 H), 1.31 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>14</sub>FN+H: 180, observed: 180.

**Example 34: (R,S) 7-Fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 26, (R,S) 7-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 3-fluorophenethylamine as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.09 (dd, J=6, 8 Hz, 1 H), 6.85-6.78 (m, 2 H), 3.10-2.89 (m, 6 H), 2.71 (dd, J=7, 13 Hz, 1 H), 1.91 (bs, 1 H), 1.33 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>14</sub>FN+H: 180, observed: 180.

**Example 35: (R,S) 7-Chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 26, (R,S) 7-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 3-chlorophenethylamine as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.10 (d, J=8 Hz, 1 H), 7.06 (m, 2 H), 3.1-2.9 (m, 6 H), 2.70 (dd, J=13, 7 Hz, 1 H), 1.89 (bs, 1 H), 1.31 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>14</sub>ClN+H: 196, observed: 196.

**Example 36: (R,S) 7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 26, (R,S) 7,8-dichloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 3,4-dichlorophenethylamine as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.20 (s, 1 H), 7.16 (s, 1 H), 3.05-2.86 (m, 6 H), 2.71 (dd, J=7, 13 Hz, 1 H), 1.83 (bs, 1 H), 1.33 (d, J=7 Hz, 3 H). MS calculated for C<sub>11</sub>H<sub>13</sub>Cl<sub>2</sub>N+H: 230, observed: 230.

**Example 37: (R,S) N-Methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




5 By the same general procedure as in example 9, (R,S) N-methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from (R,S) 8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine as a colorless oil. MS calculated for  $C_{12}H_{16}ClN+H$ : 210, observed: 210.

10 **Example 38: (R,S) 1-Methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



15 By the same general procedure as in example 26, (R,S) 1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 3-trifluoromethoxyphenethylamine as a colorless oil.  $^1H$  NMR (400 MHz,  $CD_3OD$ ) 7.39 (d,  $J=8$  Hz, 1 H), 7.19 (m, 1 H), 3.46 (m, 2 H), 3.38 (d,  $J=13$  Hz, 1 H), 3.29 (m, 1 H), 3.16 (m, 2 H), 3.05 (dd,  $J=13, 9$  Hz, 1 H), 1.50 (d,  $J=8$  Hz, 3 H). MS calculated for  $C_{12}H_{14}F_3NO+H$ : 246, observed: 246.


20 **Example 39: (R,S) 8-Iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**



25 By the same general procedure as in example 3, (R,S) 8-iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from N-trifluoroacetyl-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1H-3-benzazepine as a colorless oil.  $^1H$  NMR (400 MHz,  $CD_3OD$ ) d 7.79 (s, 1 H), 7.25 (s, 1 H),


3.46-3.40 (m, 3 H), 3.28-3.12 (m, 3 H), 3.07 (dd,  $J=13, 9$  Hz, 1 H), 1.47 (d,  $J=7$  Hz, 3 H). MS calculated for  $C_{12}H_{14}F_3INO+H$ : 372, observed: 372.

**Example 40: (R,S) N-Propyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 10, (R,S) N-Propyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from (R,S) 8-iodo-7-methoxy-1-methyl-1,2,4,5-tetrahydro-3H-3-benzazepine as a colorless oil. MS calculated for  $C_{15}H_{22}INO+H$ : 360, observed: 360.

**Example 41: (R,S) 1-Ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 19, (R,S) 1-ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from N-trifluoroacetyl-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine as a colorless oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.47 (s, 1 H), 6.54 (s, 1 H), 3.86 (s, 3 H), 3.20-2.97 (m, 4 H), 2.93-2.75 (m, 3 H), 2.64 (m, 1 H), 1.78 (m, 2 H), 0.95 (dd,  $J=8, 8$  Hz, 3 H). MS calculated for  $C_{13}H_{18}INO+H$ : 332, observed: 332.

**Example 42: (R,S) 7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




By the same general procedure as in example 29, (R,S) 7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine was obtained from N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine, *O*-trifluoromethanesulfonate as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.37 (dd, J=7, 7 Hz, 1 H), 7.30 (m, 2 H), 7.21 (d, J=7 Hz, 1 H), 7.14 (d, J=7 Hz, 1 H), 7.09 (s, 1 H), 6.86 (d, J=8 Hz, 1 H), 3.85 (s, 3 H), 3.2-2.9 (m, 6 H), 2.80 (m, 1 H), 2.64 (bs, 1 H), 1.38 (d, J=7 Hz, 3 H). MS calculated for C<sub>18</sub>H<sub>21</sub>NO+H: 268, observed: 268.

**Example 43: (R,S) 7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine**



By the same general procedure as in example 29, (R,S) 7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine was obtained from N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine, *O*-trifluoromethanesulfonate as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.35-7.10 (m, 5 H), 6.95 (dd, J=7, 8 Hz, 1 H), 3.2-2.9 (m, 6 H), 2.79 (dd, J=8, 13 Hz, 1 H), 2.70 (bs, 1 H), 1.38 (d, J=8 Hz, 3 H). MS calculated for C<sub>17</sub>H<sub>17</sub>F<sub>2</sub>N+H: 274, observed: 274.

**Example 44: (R,S) 7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine**




By the same general procedure as in example 29, (R,S) 7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine was obtained from N-trifluoroacetyl-8-chloro-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine




as a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.35-7.23 (m, 3 H), 7.19-7.09 (m, 2 H), 7.03 (s, 1 H), 3.15-2.85 (m, 7 H), 2.76 (dd,  $J=8, 13$  Hz, 1 H), 1.36 (d,  $J=8$  Hz, 3 H). MS calculated for  $\text{C}_{17}\text{H}_{17}\text{ClFN}+\text{H}$ : 290, observed: 290.

5      **Example 45: (R,S) 7-(2-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



By the same general procedure as in example 29, (R,S) 7-(2-trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was  
10      obtained from N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, *O*-trifluoromethanesulfonate as a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.71 (d,  $J=8$  Hz, 1 H), 7.52 (dd,  $J=7, 8$  Hz, 1 H), 7.42 (dd,  $J=7, 8$  Hz, 1 H), 7.31 (d,  $J=7$  Hz, 1 H), 7.17 (d,  $J=8$  Hz, 1 H), 7.11 (d,  $J=8$  Hz, 1 H), 3.15 (m, 1 H), 3.1-2.9 (m, 5 H), 2.76 (dd,  $J=8, 13$  Hz, 1 H), 2.37 (bs, 1 H), 1.38 (d,  $J=8$  Hz, 3 H). MS calculated for  $\text{C}_{18}\text{H}_{18}\text{F}_3\text{N}+\text{H}$ : 306, observed: 306.  
15


**Example 46: (R,S) 7-(3-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



20      By the same general procedure as in example 29, (R,S) 7-(3-trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, *O*-trifluoromethanesulfonate as a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.80 (s, 1 H), 7.73 (d,  $J=8$  Hz, 1 H), 7.57-7.48 (m, 2 H), 7.38 (d,  $J=8$  Hz, 1 H), 7.30 (s, 1 H), 7.24 (d,  $J=7$  Hz, 1 H), 3.16 (m, 1 H), 3.1-2.9 (m, 6 H), 2.79  
25


(dd, J=8, 13 Hz, 1 H), 1.38 (d, J=8 Hz, 3 H). MS calculated for  $C_{18}H_{18}F_3N+H$ : 306, observed: 306.

5 **Example 47: (R,S) 7-(4-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



By the same general procedure as in example 29, (R,S) 7-(4-trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from N-trifluoroacetyl-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine, O-trifluoromethanesulfonate as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.65 (s, 4 H), 7.38 (d, J=8 Hz, 1 H), 7.31 (s, 1 H), 7.24 (d, J=8 Hz, 1 H), 3.15 (m, 1 H), 3.1-2.9 (m, 5 H), 2.80 (dd, J=8, 13 Hz, 1 H), 2.48 (bs, 1 H), 1.38 (d, J=8 Hz, 3 H). MS calculated for  $C_{18}H_{18}F_3N+H$ : 306, observed: 306.


15 **Example 48: (R,S) 8-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



A solution of N-trifluoroacetyl-8-bromo-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (84 mg, 0.229 mmol) in dimethylformamide (2.5 mL) was treated with 2-chlorophenylboronic acid (43 mg, 0.275 mmol), CsF (52 mg, 0.34 mmol), water (70 mg, 3.9 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (27 mg, 0.023 mmol) and stirred overnight at 75 °C. The product mixture was diluted with EtOAc (20 mL), washed with water (10 mL), brine (10 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (10-20% EtOAc in hexane, silica) resulted in 36 mg of a clear oil. MS calculated for  $C_{19}H_{17}ClF_3NO+H$ : 368, observed: 368. The product (39 mg, 0.106 mmol) was dissolved in methanol (2 mL) treated with 15% aqueous NaOH

(2 mL), and stirred overnight at 20 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and the solvent evaporated to give 18 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.44 (d, J=8 Hz, 1 H), 7.35-7.17 (m, 5 H), 7.12 (d, J=8 Hz, 1 H), 3.14 (m, 1 H), 3.1-2.9 (m, 5 H), 2.79 (dd, J=7, 13 Hz, 1 H), 2.36 (bs, 1 H), 1.36 (d, J=7 Hz, 3 H). MS calculated for C<sub>17</sub>H<sub>18</sub>ClN<sub>3</sub>+H: 272, observed: 272.

**Example 49: (R,S) 7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




10

A solution of N-trifluoromethylacetyl-8-iodo-7-methoxy-1-methyl-1,2,4,5-tetrahydro-3H-3-benzazepine (135 mg, 0.327 mmol) in dimethylformamide (3 mL) and toluene (0.5 mL) was treated with sodium trifluoroacetate (133 mg, 0.981 mmol), copper (I) iodide (124 mg, 0.654 mmol) and the toluene distilled off to remove any residual water. The reaction mixture was stirred at 155 C for 3.5 hours, diluted with EtOAc, filtered, absorbed on silica and purified by flash chromatography (10% EtOAc in hexane, silica) resulting in 26 mg of a colorless oil. MS calculated for C<sub>15</sub>H<sub>15</sub>F<sub>6</sub>NO<sub>2</sub>+H: 356, observed: 356. The intermediate (26 mg, 0.073 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred 0.5 hours at 50 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 14 mg of a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.32 (s, 1 H), 6.73 (s, 1 H), 3.89 (s, 3 H), 3.1-2.9 (bm, 6 H), 2.75 (bm, 1 H), 2.23 (bs, 1 H), 1.36 (d, J=8 Hz, 3 H). MS calculated for C<sub>13</sub>H<sub>16</sub>F<sub>3</sub>NO+H: 260, observed: 260.

20


25

**Example 50: (R,S) 7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1H-3-benzazepine**




A solution of N-trifluoromethylacetyl-8-iodo-7-methoxy-1-methyl-1,2,4,5-tetrahydro-3H-3-benzazepine (100 mg, 0.242 mmol) in dimethylformamide (3 mL) and toluene (1 mL) was treated with sodium pentafluoropropionate (64 mg, 0.344 mmol), copper (I) iodide (92 mg, 0.484 mmol) and the toluene distilled off to remove any residual water. The reaction mixture was stirred at 160 C for 3.5 hours, diluted with EtOAc, filtered, absorbed on silica and purified by flash chromatography (10% EtOAc in hexane, silica) resulting in 22 mg of a colorless oil. MS calculated for  $C_{16}H_{15}F_8NO_2 + H$ : 406, observed: 406. The intermediate (22 mg, 0.054 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred 0.5 hours at 50 C. The product mixture was diluted with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with  $Na_2SO_4$  and concentrated to give 14 mg of a colorless oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.25 (s, 1 H), 6.74 (s, 1 H), 3.85 (s, 3 H), 3.1-2.9 (bm, 6 H), 2.76 (bm, 1 H), 2.37 (bs, 1 H), 1.35 (d,  $J=8$  Hz, 3 H). MS calculated for  $C_{14}H_{16}F_5NO + H$ : 310, observed: 310.

**Example 51: (R,S) 8-Trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



By the same general procedure as in example 26, (R,S) 8-trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine was obtained from 4-trifluoromethylphenethylamine as a colorless oil.  $^1H$  NMR (400 MHz, DMSO)  $\delta$  7.55 (d,  $J=8$  Hz, 1 H), 7.49 (s, 1 H), 7.43 (d,  $J=8$  Hz, 1 H), 3.55-3.50 (m, 1H) 3.43-3.23 (m, 7 H), 3.13 (dd,  $J=16, 7$  Hz, 1H), 3.0-2.91 (m, 2H), 1.36 (d,  $J=7$  Hz, 3 H). MS calculated for  $C_{12}H_{14}F_3N + H$ : 230.19, observed: 230.4

**Example 52: (R,S) 8-bromo-1-methoxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine**




5 A solution of 8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (0.075 g, 0.26 mmol) in dichloromethane (2 mL) was treated with BOC<sub>2</sub>O (0.062 g, 0.29 mmol), and stirred overnight at 20 C. The product was absorbed on silica and purified by flash chromatography (33% EtOAc in hexane, silica) resulting in 0.034 g of a clear oil. MS calculated for

10 C<sub>17</sub>H<sub>24</sub>BrNO<sub>4</sub>+H: 386, observed: 386. The BOC-protected intermediate was dissolved in dimethylformamide (1 mL), treated with excess NAH and excess iodomethane sequentially, and then stirred for 1 hour at 20 C. The reaction mixture was quenched with water (5 mL), extracted twice with EtOAc (5 mL), the combined organic phases were washed with brine (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and

15 concentrated to give 0.019 g of a clear oil. MS calculated for C<sub>18</sub>H<sub>26</sub>BrNO<sub>4</sub>+H: 400, observed: 400. The N-BOC protected methylether was then treated with 4M HCl in dioxane (1 mL) and stirred 2 hours at 20 C. Evaporation resulted in 0.009 g of the desired product as a clear oil. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.30 (s, 1 H), 6.92 (s, 1 H), 3.87 (s, 3H), 3.65 (s, 3H) 3.5-3.1 (m, 9 H). MS calculated for

20 C<sub>13</sub>H<sub>18</sub>BrNO<sub>2</sub>+H: 300, observed: 300.

**Example 53: (R,S) 8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Crotyl, *N*-trifluoroacetyl-2-iodo-4-chlorophenethylamine

25 A solution of *N*-trifluoroacetyl-2-iodo-4-chlorophenethylamine (6.2 g, 15.8 mmol) in dimethylformamide (350 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (15.8 g, 114 mmol) and crotyl bromide (6.0 g, 44 mmol) sequentially, the mixture was stirred

at 60 C for 16 hours and then cooled to 20 C. The mixture was diluted with EtOAc (350 mL), washed with water (3 x 300 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (5-15% EtOAc in hexane) resulted in 2.5 g of a clear oil. MS calculated for C<sub>14</sub>H<sub>14</sub>ClF<sub>3</sub>INO+H: 432, observed: 432.

5

*N-Trifluoroacetyl-8-chloro-1-ethyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-crotyl, N-trifluoroacetyl-2-iodo-4-chlorophenethylamine (2.5 g, 5.8 mmol) in dimethylformamide (250 mL) was treated with KOAc (1.07 g, 10.9 mmol), n-Bn<sub>2</sub>Et<sub>2</sub>NBr (1.33 g, 5.84 mmol), Pd(OAc)<sub>2</sub> (0.063 g, 0.28 mmol) and stirred overnight at 77 C. The product mixture was cooled to 20 C, filtered, diluted with water (100 mL), extracted with EtOAc (3 x 100 mL), the combined organic phases washed with water (100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. Flash chromatography (2-20% EtOAc in hexane, silica) resulted in 0.339 g of a clear oil. The product, which was assumed to be a mixture of double-bond isomers, was dissolved in methanol (50 mL) treated with Et<sub>3</sub>N (0.2 mL), 10% Pd/C (0.10 g) and stirred 16 hours under 100 psi of hydrogen. The product mixture was filtered, concentrated and purified by flash chromatography (5% EtOAc in hexane, silica) resulting in 0.20 g of a white solid. MS calculated for C<sub>14</sub>H<sub>15</sub>ClF<sub>3</sub>NO+H: 306, observed: 306.


15  
20

*8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of N-trifluoroacetyl-8-chloro-1-ethyl-2,3,4,5-tetrahydro-1H-3-benzazepine (63 mg, 0.207 mmol) in methanol (2 mL) was treated with 15% aqueous NaOH (2 mL), and stirred for 3.5 hours at 60 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 35 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.2 (m, 3 H), 3.3-3.0 (m, 7 H), 1.9-1.6 (m, 2 H), 0.91 (t, J=7 Hz, 3 H). MS calculated for C<sub>12</sub>H<sub>16</sub>ClN+H: 210, observed: 210.

25

**Example 54: (R,S) 8-Chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine**



*N*-Trifluoroacetyl-8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine

5 A solution of *N*-trifluoroacetyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (2.5 g, 8.5 mmol) in 1,2-dichloroethane (15 mL) was treated with Selectfluor (3.9 g, 11 mmol), trifluoromethanesulfonic acid (8 mL, 90 mmol) and stirred 60 hours at 75 C. The product mixture was poured into water (200 mL), extracted with EtOAc (200 mL), the organic phase washed with saturated aqueous

10 NaHCO<sub>3</sub> (2 x 100 mL), brine (100 mL), dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography (6 % EtOAc in hexane, silica) resulting in 1.6 g of a white solid. MS calculated for C<sub>13</sub>H<sub>12</sub>ClF<sub>4</sub>NO+H: 310, observed: 310.

15 *8-Chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine*

A solution of *N*-trifluoroacetyl-8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine (160 mg, 0.22 mmol) in methanol (3 mL) was treated with 15% aqueous NaOH (2 mL), and stirred for 3.5 hours at 25 C. The product mixture was concentrated, extracted 3 times with CH<sub>2</sub>Cl<sub>2</sub> (5 mL), dried

20 with Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 93 mg of a clear oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.11 (m, 1 H), 6.85 (m, 1 H), 3.05-2.95 (m, 3 H), 2.95-2.80 (m, 3H), 2.68 (m, 1 H), 2.38 (bm, 1 H), 1.31 (m, 3 H). MS calculated for C<sub>11</sub>H<sub>13</sub>ClFN+H: 214, observed: 214.

25 **Example 55 Separation of enantiomers for selected compounds of the invention**

The following compounds were separated into their respective enantiomers using a Varian ProStar HPLC system with a 20 mm x 250 mm Chiralcel OD chiral column, eluting with 0.2 % diethylamine in various concentrations of isopropanol (IPA) in hexanes, see Table 1 below. In some cases, the separations were

30 performed on the intermediate trifluoroacetamide protected amines.

Table 1

Example	Enantiomer	Retention time for the free amine (mins)	Retention time for the trifluoroacetamide	Conditions
1	1	21.9		5% IPA in hexane
	2	24.5		10 mL/min
2	1	42		5% IPA in hexane
	2	47		9 mL/min
3	1	20.8		5% IPA in hexane
	2	24.2		10 mL/min
19	1	34.9		1% IPA in hexane
	2	39.5		9 mL/min
26	1		23.8 <sup>1</sup>	5% IPA in hexane
	2		29.2 <sup>2</sup>	7 mL/min
37	1		23.8 <sup>3</sup>	5% IPA in hexane
	2		29.2 <sup>4</sup>	7 mL/min
51	1		18.6 <sup>5</sup>	1% IPA in hexane
	2		21.4 <sup>6</sup>	9 mL/min
53	1		13.7 <sup>7</sup>	5% IPA in hexane
	2		20.2 <sup>8</sup>	10 mL/min

<sup>1</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 1 of Compound 26.

<sup>2</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 2 of Compound 26.

<sup>3</sup> The separated trifluoroacetamide enantiomer was hydrolyzed and subsequently N-methylated to give Enantiomer 1 of Compound 37.

<sup>4</sup> The separated trifluoroacetamide enantiomer was hydrolyzed and subsequently N-methylated to give Enantiomer 2 of Compound 37.

<sup>5</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 1 of Compound 51.

<sup>6</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 2 of Compound 51.

<sup>7</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 1 of Compound 53.

<sup>8</sup> The separated trifluoroacetamide enantiomer was hydrolyzed to give Enantiomer 2 of Compound 53.



**Example 56****Intracellular IP<sub>3</sub> Accumulation Assay**

HEK293 cells were transfected in 15cm sterile dishes with or without (control) 16ug of human 5-HT<sub>2C</sub> receptor cDNA using 25ul of lipofectamine. Cells were then incubated for 3-4 hours at 37°C/5%CO<sub>2</sub> and then transfection media was removed and replaced with 100ul of DMEM. Cells were then plated onto 100cm sterile dishes. The next day cells were plated into 96 well PDL microtiter plates at a density of 55K/0.2ml. Six hours latter, media was exchanged with [<sup>3</sup>H]inositol (0.25 uCi/well) in inositol free DMEM and plates were incubated at 37°C/5%CO<sub>2</sub> overnight. The next day, wells were aspirated and 200ul of DMEM containing test compound, 10uM pargyline, and 10mM LiCl was added to appropriate wells. Plates were then incubated at 37°C/5%CO<sub>2</sub> for three hours followed aspiration and by addition of fresh ice cold stop solution (1M KOH, 19mM Na-borate, 3.8 mM EDTA) to each well. Plates were kept on ice for 5-10 min and the wells were neutralized by addition of 200ul of fresh ice cold neutralization solution (7.5% HCl). Plates were then frozen until further processing is desired. The lysate was then transferred into 1.5 ml Eppendorf tubes and 1 ml of chloroform/methanol (1:2) was added/tube. The solution was vortexed for 15 seconds and the upper phase was applied to a Biorad AG1-X8™ anion exchange resin (100-200 mesh). First, the resin was washed with water at 1:1.25 W/V and 0.9 ml of upper phase was loaded onto the column. The column was then washed with 10 ml of 5 mM myo-inositol and 10 ml of 5 mM Na-borate/60mM Na-formate. The inositol tris phosphates were eluted into scintillation vials containing 10 ml of scintillation cocktail with 2 ml of 0.1 M formic acid/ 1 M ammonium formate. The columns were regenerated by washing with 10 ml of 0.1 M formic acid/3M ammonium formate and rinsed twice with dd H<sub>2</sub>O and stored at 4°C in water.

The biological activities in the IP Accumulation Assay for several representative compounds are shown in Table 2 below:

Table 2

Compound (Example Number)	5-HT <sub>2C</sub> (IC <sub>50</sub> )* IP Accumulation Assay (nM)
1	4.2
2	4.5
3	1.4
4	2.1
5	12.1
12	6.3
19	18
26	5.8
32	2.1

\* Reported values are averages of at least two trials.

The majority of the other compounds of the Examples were tested at least once, and they showed activities in the IP Accumulation Assay in the range between ~1.4 nM and ~5 ?M.

### Example 57

#### Inhibition of food intake in food-deprived rats

Male Sprague-Dawley rats (250-350g) were deprived of food overnight prior to testing. Prior to food deprivation, the animals were weighed and separated into treatment groups in order to balance groups according to body weight. On the test day, animals were placed into individual cages (no bedding) at 9:00am with free access to water. At 10:00am, animals were injected with test compound (p.o., i.p., or s.c.) and then presented with a pre-weighed amount of food in a dish either 60min (p.o.) or 30min (i.p. and s.c.) after drug administration. Food consumption over different time points was then determined by weighing the food cup at 1, 2, 4, and 6hr after the food was presented. Thus, food consumption was measured at 2, 3, 5, and 7hr post-injection in p.o. studies, and at 1.5, 2.5, 4.5, and 6.5hr post-injection in i.p. and s.c. studies.


Figures 1A-G illustrate the effects of seven different compounds on food intake in food-deprived rats. All compounds inhibited food intake dose-dependently. This effect was consistently most pronounced over the first 1hr after food presentation. Some compounds (Figures 1A, 1C, and 1E) maintained an inhibitory effect on food intake relative to vehicle-treated controls at 6hr after food presentation. Compounds were also shown to be effective via all routes of administration including p.o.

It is intended that each of the patents, applications, printed publications, and other published documents mentioned or referred to in this specification be herein incorporated by reference in their entirety.

Those skilled in the art will appreciate that numerous changes and modifications can be made to the preferred embodiments of the invention and that such changes and modifications can be made without departing from the spirit of the invention. It is therefore intended that the appended claims cover all such equivalent variations as fall within the true spirit and scope of the invention.

**What is claimed is:**

1. A compound of Formula (I):



wherein:

$R_1$  is H or  $C_{1-8}$  alkyl;

$R_2$  is  $C_{1-8}$  alkyl,  $-CH_2-O-C_{1-8}$  alkyl,  $-C(=O)-O-C_{1-8}$  alkyl,  $-C(=O)-NH-C_{1-8}$  alkyl, OH, or  $CH_2OH$ ;

$R_{2a}$  is H;

or  $R_2$  and  $R_{2a}$  together form  $-CH_2-CH_2-$ ;

$R_3$  and  $R_4$  are each independently H, halogen, perhaloalkyl, CN,  $OR_5$ ,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhalo alkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

or  $R_3$  and  $R_4$  together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each  $R_5$  is independently  $C_{1-8}$  alkyl,  $C_{1-8}$  alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

$R_6$  is H or  $C_{1-8}$  alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(A) if  $R_2$  is methyl and  $R_1$  and  $R_3$  are both H, then  $R_4$  is not thiazole, substituted thiazole or a thiazole derivative:

(B) if  $R_6$  is other than H, then neither  $R_3$  nor  $R_4$  can be H;

(C) if  $R_1$  and  $R_2$  are methyl, and  $R_4$  is H, then  $R_3$  cannot be  $NHR_5$  or  $N(R_5)_2$ ;

(D) if  $R_1$  and  $R_2$  are methyl, and  $R_4$  is H, then  $R_3$  cannot be imidazole, substituted imidazole, or an imidazole derivative; and

5 (E) if  $R_3$  is OH, and  $R_1$  is methyl then  $R_2$  cannot be cyclopentyl,  $-CH_2$ -cyclohexyl, cyclopropylmethyl, or cyclohexyl.

2. The compound according to claim 1 wherein  $R_1$  is H.

10 3. The compound according to claim 1 wherein  $R_1$  is methyl.

4. The compound according to claim 1 wherein  $R_2$  is methyl, ethyl, n-propyl or isopropyl.

15 5. The compound according to claim 1 wherein  $R_2$  is methyl or ethyl.

6. The compound according to claim 1 wherein  $R_2$  and  $R_{2a}$  together form  
 $-CH_2-CH_2-$ .

20

7. The compound according to claim 1 wherein  $R_3$  is chlorine.

8. The compound according to claim 1 wherein  $R_3$  is bromine.

25 9. The compound according to claim 1 wherein  $R_3$  is perhaloalkyl.

10. The compound according to claim 9 wherein  $R_3$  is  $CF_3$ .

30 11. The compound according to claim 1 wherein  $R_3$  is selected from the group consisting of thiophenyl, furanyl, pyrrolyl, pyrazolyl and imidazolyl.

12. The compound according to claim 1 wherein R<sub>4</sub> is methoxy,  
ethoxy,  
n-propoxy, isopropoxy or allyloxy.
- 5
13. The compound according to claim 1 wherein R<sub>4</sub> is selected from  
the group consisting of thiophenyl, furanyl, pyrrolyl, pyrazolyl and  
imidazolyl optionally substituted with one or two substituents  
selected from halogen or methyl.
- 10
14. The compound according to claim 1 wherein R<sub>4</sub> is phenyl  
optionally substituted with up to two substituents selected from C<sub>1-8</sub>  
alkyl, halogen, and alkoxy.
- 15
15. The compound according to claim 1 wherein R<sub>3</sub> and R<sub>4</sub> taken  
together form -O-CH=C(CH<sub>3</sub>)-.
16. The compound according to claim 1 wherein R<sub>3</sub> is halogen and R<sub>4</sub>  
is -OR<sub>5</sub> wherein R<sub>5</sub> is C<sub>1-8</sub> alkyl.
- 20
17. The compound according to claim 1 wherein R<sub>3</sub> is chlorine or  
bromine and R<sub>4</sub> is methoxy.
18. The compound according to claim 1 wherein R<sub>3</sub> is halogen and R<sub>4</sub>  
is allyloxy.
- 25
19. The compound according to claim 1 wherein R<sub>3</sub> is H and R<sub>4</sub> is 5-  
membered heteroaryl ring having up to two heteroatoms selected  
from O, N and S, and up to two substituents selected from halogen  
and C<sub>1-8</sub> alkyl, or R<sub>4</sub> is phenyl optionally substituted with up to two  
substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy.
- 30

20. The compound according to claim 1 wherein  $R_3$  is H and  $R_4$  is a disubstituted pyrrazole or monohalo-substituted phenyl.

5 21. The compound according to claim 1 wherein said substituents of said pyrrazole are bromine and methyl.

22. The compound according to claim 1 wherein:

10  $R_2$  is methyl, ethyl, isopropyl, or  $CH_2OH$ ; or  $R_2$  and  $R_{2a}$  taken together form  $-CH_2-CH_2-$ ;

$R_3$  is H, halogen, or a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S, and up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

15  $R_4$  is H, alkoxy, a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S and up to two substituents selected from halogen and  $C_{1-8}$  alkyl, or phenyl optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, and alkoxy;

or  $R_3$  and  $R_4$  taken together form  $-O-CH=C(CH_3)-$ ; and

20  $R_6$  is H or methyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

23. The compound according to claim 1 wherein:

25  $R_2$  is methyl, ethyl, isopropyl, or  $CH_2OH$ ; or  $R_2$  and  $R_{2a}$  taken together form  $-CH_2-CH_2-$ ;

$R_3$  is chlorine, bromine, or iodine;

$R_4$  is alkoxy; and

$R_6$  is H or methyl or a pharmaceutically acceptable salt, solvate or hydrate thereof.

30

24. The compound according to claim 1 wherein:

R<sub>1</sub> is H;

R<sub>2</sub> is methyl;

R<sub>3</sub> is H, chlorine, bromine, or thiophene;

R<sub>4</sub> is alkoxy, pyrrazoly-3-yl or phenyl wherein said  
pyrrazole optionally has up to two substituents selected from  
halogen and C<sub>1-8</sub> alkyl, and said phenyl optionally has a single  
halogen substituent; and

R<sub>6</sub> is H or a pharmaceutically acceptable salt, solvate or  
hydrate thereof.

25. A compound of claim 1 selected from the group consisting of:
- 8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 7-Benzoyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- N-Propyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 7-Hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 3,5-Dimethyl-6,7,8,9-tetrahydro-5*H*-1-oxa-7-aza-cycloheptaindene;
- 7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;
- 8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-



- benzazepine;  
8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
5 8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
10 8-Bromo-7-methoxy-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
15 7-(2-Methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(4-Bromo-2-methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(3-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
20 8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
25 N-Methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
1-Methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Propyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
30 1-Ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

- 7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
5 7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
10 7-(3-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(4-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
and  
15 8-bromo-1-methoxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

26. A compound according to claim 1 selected from the group  
20 consisting of:  
8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
25 8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and  
30

7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

- 5                    27.    A compound according to claim 1 selected from the group consisting of:
- 8-Chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
10                   8-Trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
15                   7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7,8-Dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
and  
8-Chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or  
20                   a pharmaceutically acceptable salt, solvate or hydrate thereof.
- 25                   28.    A pharmaceutical composition comprising a compound according to any one of claims 1, 25, 26 and 27 and a pharmaceutically acceptable carrier or excipient.
29.    A method of modulating a 5HT<sub>2C</sub> receptor comprising contacting said receptor with a pharmaceutically effective amount of a compound according to any one of claims 1, 25, 26 and 27.
- 30                   30.    The method of claim 29 wherein said compound is an agonist of said receptor.

31. A method of prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea comprising administering to a patient in need of such prophylaxis or treatment an effective dose of a compound as in any one of claims 1, 25, 26 and 27.
32. A method according to claim 31 wherein the disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.
33. A method of decreasing food intake of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 25, 26 and 27.
34. A method of inducing satiety in a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 25, 26 and 27.

35. A method of controlling weight gain of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 25, 26 and 27.
- 5 36. A method of prophylaxis or treatment of obesity comprising administering to a patient in need of such prophylaxis or treatment a pharmaceutically effective amount of a compound as in any one of claims 1, 25, 26 and 27.
- 10 37. The method of claim 36 further comprising the step of identifying a subject, said subject being in need of prophylaxis or treatment for obesity, wherein said identifying step is performed prior to administering to said subject said pharmaceutically effective amount of said compound.

1/7




Fig. 1A

2/7




Fig. 1B

3/7




Fig. 1C



4/7




Fig. 1D

5/7




Fig. 1E

6/7




Fig. 1F

7/7




Fig. 1G

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number  
**WO 2003/086306 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 223/14**,  
223/16, 487/00, 491/00, 513/00, A61K 31/55

(74) Agents: **STRAHER, Michael, P.** et al.; Cozen O'Connor,  
1900 Market Street, Philadelphia, PA 19103 (US).

(21) International Application Number:  
PCT/US2003/011076

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 11 April 2003 (11.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/372,058 12 April 2002 (12.04.2002) US  
60/405,495 23 August 2002 (23.08.2002) US  
60/434,607 18 December 2002 (18.12.2002) US  
10/410,991 10 April 2003 (10.04.2003) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ARENA PHARMACEUTICALS, INC.** [US/US]; 6166 Nancy Ridge Drive, San Diego, CA 92121 (US).

Published:  
— *with international search report*

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **SMITH, Brian** [US/US]; 11184 Vista Sorrento Parkway, Apartment G-301, San Diego, CA 92130 (US). **SMITH, Jeffrey** [US/US]; 8041 Jade Coast Road, San Diego, CA 92126 (US).

(88) Date of publication of the international search report:  
19 February 2004

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 5HT<sub>2C</sub> RECEPTOR MODULATORS

(57) Abstract: The present invention relates to novel compounds of: (Formula I); which act as 5HT<sub>2C</sub> receptor modulators. These compounds are useful in pharmaceutical compositions whose use includes the treatment of obesity.



WO 2003/086306 A3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/11076

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 223/14, 223/16, 487/00, 491/00, 513/00; A61K 31/55  
US CL : 540/543, 586, 593; 514/212.02, 215, 217.01

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/543, 586, 593; 514/212.02, 215, 217.01

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3,652,543 (HOEGERLE et al.) 28 March 1972 (28.03.1972), column 8, lines 25-35.	1-27
X	US 4,111,957 (HOLDEN et al.) 05 September 1978 (05.09.1978), column 8, lines 26-27.	1-27
X	US 4,988,690 (EFFLAND et al.) 29 January 1991 (29.01.1991), column 12, lines 31-32.	1-27
Y	US 5,015,639 (BERGER et al.) 14 May 1991 (14.05.1991), columns 1-2.	1-37

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

25 September 2003 (25.09.2003)

Date of mailing of the international search report

16 OCT 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Bruck Korte, Ph.D.

Telephone No. 703-308-1235

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
23 October 2003 (23.10.2003)

PCT

(10) International Publication Number  
**WO 2003/086306 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 223/14**,  
223/16, 487/00, 491/00, 513/00, A61K 31/55

(21) International Application Number:  
PCT/US2003/011076

(22) International Filing Date: 11 April 2003 (11.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/372,058 12 April 2002 (12.04.2002) US  
60/405,495 23 August 2002 (23.08.2002) US  
60/434,607 18 December 2002 (18.12.2002) US  
10/410,991 10 April 2003 (10.04.2003) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): ARENA PHARMACEUTICALS, INC. [US/US]; 6166 Nancy Ridge Drive, San Diego, CA 92121 (US).

Published:  
— with international search report  
— with amended claims

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SMITH, Brian [US/US]; 11184 Vista Sorrento Parkway, Apartment G-301, San Diego, CA 92130 (US). SMITH, Jeffrey [US/US]; 8041 Jade Coast Road, San Diego, CA 92126 (US).


(88) Date of publication of the international search report:  
19 February 2004

Date of publication of the amended claims: 25 March 2004

(74) Agents: STRAHER, Michael, P. et al.; Cozen O'Connor, 1900 Market Street, Philadelphia, PA 19103 (US).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 5HT<sub>2C</sub> RECEPTOR MODULATORS



(57) Abstract: The present invention relates to novel compounds of: (Formula I); which act as 5HT<sub>2C</sub> receptor modulators. These compounds are useful in pharmaceutical compositions whose use includes the treatment of obesity.


WO 2003/086306 A3

## AMENDED CLAIMS

[Received by the International Bureau on 21 November 2003 (21.11.03):  
claims 2-10, 14, 15, 30, 32 and 37 are unchanged. Original claims 12, 16-21, 23 and 27 are canceled.  
Original claims 1, 11, 13, 22, 24-26, 28, 29, 31 and 33-36 are renumbered.  
New claims 38-116 are added.]

**What is claimed is:**

1. A compound of Formula (I):



wherein:

$R_1$  is H or  $C_{1-8}$  alkyl;

$R_2$  is  $C_{1-8}$  alkyl,  $-CH_2-O-C_{1-8}$  alkyl,  $-C(=O)-O-C_{1-8}$  alkyl,  $-C(=O)-NH-C_{1-8}$  alkyl, OH, or  $CH_2OH$ ;

$R_{2a}$  is H;

or  $R_2$  and  $R_{2a}$  together form  $-CH_2-CH_2-$ ;

$R_3$  is halogen, perhaloalkyl, CN,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

$R_4$  is H, halogen, perhaloalkyl, CN,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

or  $R_3$  and  $R_4$  together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each  $R_5$  is independently  $C_{1-8}$  alkyl,  $C_{1-8}$  alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

$R_6$  is H or  $C_{1-8}$  alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(B) if  $R_6$  is other than H, then neither  $R_3$  nor  $R_4$  can be H;



(C) if  $R_1$  and  $R_2$  are methyl, and  $R_4$  is H, then  $R_3$  cannot be  $NHR_5$  or  $N(R_5)_2$ ; and

(D) if  $R_1$  and  $R_2$  are methyl, and  $R_4$  is H, then  $R_3$  cannot be imidazole, substituted imidazole, or an imidazole derivative.

2. The compound according to claim 1 wherein  $R_1$  is H.
3. The compound according to claim 1 wherein  $R_1$  is methyl.
4. The compound according to claim 1 wherein  $R_2$  is methyl, ethyl, n-propyl or isopropyl.
5. The compound according to claim 1 wherein  $R_2$  is methyl or ethyl.
6. The compound according to claim 1 wherein  $R_2$  and  $R_{2a}$  together form  $-CH_2-CH_2-$ .
7. The compound according to claim 1 wherein  $R_3$  is chlorine.
8. The compound according to claim 1 wherein  $R_3$  is bromine.
9. The compound according to claim 1 wherein  $R_3$  is perhaloalkyl.
10. The compound according to claim 9 wherein  $R_3$  is  $CF_3$ .
11. The compound according to claim 1 wherein  $R_3$  is selected from the group consisting of thienyl, furanyl, pyrrolyl, pyrazolyl and imidazolyl.
13. The compound according to claim 1 wherein  $R_4$  is selected from the group consisting of thienyl, furanyl, pyrrolyl, pyrazolyl and imidazolyl optionally substituted with one or two substituents selected from halogen or methyl.

14. The compound according to claim 1 wherein R<sub>4</sub> is phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy.
15. The compound according to claim 1 wherein R<sub>3</sub> and R<sub>4</sub> taken together form -O-CH=C(CH<sub>3</sub>)-.
22. The compound according to claim 1 wherein:  
R<sub>2</sub> is methyl, ethyl, isopropyl, or CH<sub>2</sub>OH; or R<sub>2</sub> and R<sub>2a</sub> taken together form -CH<sub>2</sub>-CH<sub>2</sub>-;  
R<sub>3</sub> is halogen, or a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S, and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;  
R<sub>4</sub> is H, a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N and S and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, or phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy;  
or R<sub>3</sub> and R<sub>4</sub> taken together form -O-CH=C(CH<sub>3</sub>)-; and  
R<sub>6</sub> is H or methyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof.
24. The compound according to claim 1 wherein:  
R<sub>1</sub> is H;  
R<sub>2</sub> is methyl;  
R<sub>3</sub> is chlorine, bromine, or thiophene;  
R<sub>4</sub> is pyrazol-3-yl or phenyl wherein said pyrazole optionally has up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, and said phenyl optionally has a single halogen substituent; and  
R<sub>6</sub> is H or a pharmaceutically acceptable salt, solvate or hydrate thereof.
25. A compound selected from the group consisting of:

8-Bromo-7-hydroxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Benzoyloxy-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-7-ethoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-7-isopropoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Propyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Hydroxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
3,5-Dimethyl-6,7,8,9-tetrahydro-5*H*-1-oxa-7-aza-cycloheptaindene;  
7-Allyloxy-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Methoxy-1-methyl-8-(2-thienyl)-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Cyano-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-bromo-1-cyclopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-bromo-1-hydroxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-isopropyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-7-hydroxy-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-bromo-1-isopropyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-7-methoxy-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Allyloxy-8-bromo-1,4-dimethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-Methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(4-Bromo-2-methyl-2*H*-pyrazol-3-yl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(3-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-1-hydroxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

7,8-Dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Methyl-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
1-Methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-1-methyl-7-trifluoromethoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Propyl-8-iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
1-Ethyl-8-iodo-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(3-Methoxyphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2,6-difluorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-fluorophenyl)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(2-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(3-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-(4-Trifluoromethylphenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-(2-Chlorophenyl)-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; and  
8-bromo-1-methoxymethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate thereof.

26. A compound selected from the group consisting of:

8-Bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Iodo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
N-Methyl-8-bromo-7-methoxy-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Bromo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
8-Chloro-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;

8-Iodo-1-ethyl-7-methoxy-2,3,4,5-tetrahydro-1*H*-3-benzazepine;  
7-Methoxy-1-methyl-8-trifluoromethyl-2,3,4,5-tetrahydro-1*H*-3-  
benzazepine; and  
7-Methoxy-1-methyl-8-pentafluoroethyl-2,3,4,5-tetrahydro-1*H*-3-  
benzazepine; or a pharmaceutically acceptable salt, solvate or hydrate  
thereof.

28. A pharmaceutical composition comprising a compound according to any one of claims 1, 42, 51, 52, 53, and 105 and a pharmaceutically acceptable carrier or excipient.
29. A method of modulating a 5HT<sub>2C</sub> receptor comprising contacting said receptor with a pharmaceutically effective amount of a compound according to any one of claims 1, 42, 51, 52, 53, and 105.
30. The method of claim 29 wherein said compound is an agonist of said receptor.
31. A method of prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea comprising administering to a patient in need of such prophylaxis or treatment an effective dose of a compound as in any one of claims 1, 42, 51, 52, 53, and 105.
32. A method according to claim 31 wherein the disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.

33. A method of decreasing food intake of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 42, 51, 52, 53, and 105.

34. A method of inducing satiety in a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 42, 51, 52, 53, and 105.

35. A method of controlling weight gain of a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound as in any one of claims 1, 42, 51, 52, 53, and 105.

36. A method of prophylaxis or treatment of obesity comprising administering to a patient in need of such prophylaxis or treatment a pharmaceutically effective amount of a compound as in any one of claims 1, 42, 51, 52, 53, and 105.

37. The method of claim 36 further comprising the step of identifying a subject, said subject being in need of prophylaxis or treatment for obesity, wherein said identifying step is performed prior to administering to said subject said pharmaceutically effective amount of said compound.

38. The compound of claim 1 wherein:

R<sub>1</sub> is H or Me;

R<sub>2</sub> is Me, Et, or OH;

R<sub>2a</sub> is H;

R<sub>3</sub> is F, Cl, Br, I, CF<sub>3</sub>, or 2-chlorophenyl;

R<sub>4</sub> is H; and


R<sub>6</sub> is H.

39. The compound of claim 1 wherein:

$R_1$  is H;  
 $R_2$  is  $C_{1-8}$  alkyl;  
 $R_{2a}$  is H;  
 $R_3$  is halogen;  
 $R_4$  is H; and  
 $R_6$  is H.

41. A method of preparing a composition comprising combining a compound of claim 1 and a pharmaceutically acceptable carrier.

42. A compound of Formula (I):



wherein:

$R_1$  is H or straight-chained or branched  $C_{1-8}$  alkyl;  
 $R_2$  is methyl, ethyl,  $-CH_2-O-C_{1-8}$  alkyl,  $-C(=O)-O-C_{1-8}$  alkyl,  $-C(=O)-NH-C_{1-8}$  alkyl, or  $CH_2OH$ ;  
 $R_{2a}$  is H;  
 or  $R_2$  and  $R_{2a}$  together form  $-CH_2-CH_2-$ ;  
 $R_3$  is halogen, perhaloalkyl, CN,  $OR_5$ ,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;  
 $R_4$  is H, halogen, perhaloalkyl, CN,  $OR_5$ ,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhalo alkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H;

(C) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be NHR<sub>5</sub> or N(R<sub>5</sub>)<sub>2</sub>;

(D) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be imidazole, substituted imidazole, or an imidazole derivative; and

(E) if R<sub>3</sub> is OH, and R<sub>1</sub> is methyl then R<sub>2</sub> cannot be cyclopentyl, -CH<sub>2</sub>-cyclohexyl, cyclopropylmethyl, or cyclohexyl.

43. The compound of claim 42 wherein:

R<sup>1</sup> is H;

R<sup>2</sup> is methyl or ethyl;

R<sup>2a</sup> is H;

R<sup>3</sup> is F, Cl, Br, or I;

R<sup>4</sup> is H, halogen, perhaloalkyl, CN, OR<sub>5</sub>, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, OH, aryl, or heteroaryl, wherein said aryl is optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl is optionally substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

each R<sup>5</sup> is, independently, C<sub>1-8</sub> alkyl, C<sub>2-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, perhaloalkyl, or allyl; and

R<sup>6</sup> is H.



44. The compound of claim 42 wherein:

$R^1$  is H;

$R^2$  is methyl or ethyl;

$R^{2a}$  is H;

$R^3$  is F, Cl, Br, or I;

$R^4$  is H, hydroxy, methoxy, ethoxy, isopropoxy, allyloxy, benzyloxy, fluoro, chloro, trifluoromethoxy, 2-methyl-2*H*-pyrazol-3-yl, 4-bromo-2-methyl-2*H*-pyrazol-3-yl, 3-chlorophenyl, 2-chlorophenyl, 3-methoxyphenyl, 2,6-difluorophenyl, 2-fluorophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, or 4-trifluoromethylphenyl; and  
 $R^6$  is H.

45. The compound of claim 42 wherein:

$R^1$  is H;

$R^2$  is methyl or ethyl;

$R^{2a}$  is H;

$R^3$  is  $OR^5$ ;

$R^4$  is H; and

$R^6$  is H.

46. The compound of claim 42 wherein:

$R^1$  is H;

$R^2$  is methyl or ethyl;

$R^{2a}$  is H;

$R^3$  is  $OR^5$ ;


$R^4$  is H;

$R^5$  is arylalkyl or heteroarylalkyl; and

$R^6$  is H.

47. The compound according to claim 42 wherein  $R_4$  is methoxy, ethoxy, n-propoxy, isopropoxy, or allyloxy.

48. The compound according to claim 42 wherein  $R_3$  is halogen and  $R_4$  is  $-OR_5$  wherein  $R_5$  is  $C_{1-8}$  alkyl.
49. The compound according to claim 42 wherein  $R_3$  is chlorine or bromine and  $R_4$  is methoxy.
50. The compound according to claim 42 wherein  $R_3$  is halogen and  $R_4$  is allyloxy.
51. A compound of Formula (I):



wherein:

$R_1$  is H or straight-chained or branched  $C_{1-8}$  alkyl;

$R_2$  is methyl, ethyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, tert-pentyl, n-hexyl,  $-CH_2-O-C_{1-8}$  alkyl,  $-C(=O)-O-C_{1-8}$  alkyl,  $-C(=O)-NH-C_{1-8}$  alkyl, or  $CH_2OH$ ;

$R_{2a}$  is H;

or  $R_2$  and  $R_{2a}$  together form  $-CH_2-CH_2-$ ;

$R_3$  is halogen, perhaloalkyl, CN,  $OR_5$ ,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

$R_4$  is H, halogen, perhaloalkyl, CN,  $OR_5$ ,  $SR_5$ ,  $NHR_5$ ,  $N(R_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $C_{1-8}$  alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $C_{1-8}$  alkyl;

or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and


R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof  
provided that:

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H;

(C) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be NHR<sub>5</sub> or N(R<sub>5</sub>)<sub>2</sub>; and

(D) if R<sub>1</sub> and R<sub>2</sub> are methyl, and R<sub>4</sub> is H, then R<sub>3</sub> cannot be imidazole, substituted imidazole, or an imidazole derivative.

52. A compound of Formula (I):



wherein:

R<sub>1</sub> is ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, sec-butyl, tert-butyl, cyclobutyl, n-pentyl, isopentyl, tert-pentyl, cyclopentyl, n-hexyl, or cyclohexyl;

R<sub>2</sub> is C<sub>1-8</sub> alkyl, -CH<sub>2</sub>-O-C<sub>1-8</sub> alkyl, -C(=O)-O-C<sub>1-8</sub> alkyl, -C(=O)-NH-C<sub>1-8</sub> alkyl, OH, or CH<sub>2</sub>OH;

R<sub>2a</sub> is H;

or R<sub>2</sub> and R<sub>2a</sub> together form -CH<sub>2</sub>-CH<sub>2</sub>-;

R<sub>3</sub> and R<sub>4</sub> are each independently H, halogen, perhaloalkyl, CN, OR<sub>5</sub>, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, perhalo alkyl, and alkoxy, and said heteroaryl can be optionally

substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;


each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H.

53. A compound of Formula (I):



wherein:

R<sub>1</sub> is H or C<sub>1-8</sub> alkyl;

R<sub>2</sub> is C<sub>1-8</sub> alkyl, -CH<sub>2</sub>-O-C<sub>1-8</sub> alkyl, -C(=O)-O-C<sub>1-8</sub> alkyl, -C(=O)-NH-C<sub>1-8</sub> alkyl, OH, or CH<sub>2</sub>OH;

R<sub>2a</sub> is H;

or R<sub>2</sub> and R<sub>2a</sub> together form -CH<sub>2</sub>-CH<sub>2</sub>-;

R<sub>3</sub> is H, halogen, perhaloalkyl, CN, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, perhaloalkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

R<sub>4</sub> is halogen, perhaloalkyl, CN, SR<sub>5</sub>, NHR<sub>5</sub>, N(R<sub>5</sub>)<sub>2</sub>, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, perhaloalkyl, and alkoxy,

and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and C<sub>1-8</sub> alkyl;

or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each R<sub>5</sub> is independently C<sub>1-8</sub> alkyl, C<sub>1-8</sub> alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

R<sub>6</sub> is H or C<sub>1-8</sub> alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof

provided that:

(A) if R<sub>2</sub> is methyl and R<sub>1</sub> and R<sub>3</sub> are both H, then R<sub>4</sub> is not thiazole, substituted thiazole or a thiazole derivative; and

(B) if R<sub>6</sub> is other than H, then neither R<sub>3</sub> nor R<sub>4</sub> can be H.

54. The compound according to claim 53 wherein R<sub>3</sub> is H and R<sub>4</sub> is a 5-membered heteroaryl ring having up to two heteroatoms selected from O, N, and S, and up to two substituents selected from halogen and C<sub>1-8</sub> alkyl, or R<sub>4</sub> is phenyl optionally substituted with up to two substituents selected from C<sub>1-8</sub> alkyl, halogen, and alkoxy.

55. The compound according to claim 53 wherein R<sub>3</sub> is H and R<sub>4</sub> is a disubstituted pyrrazole or monohalo-substituted phenyl.

56. The compound according to claim 55 wherein said substituents of said disubstituted pyrrazole are bromine and methyl.

57. A compound according to claim 1 that is 8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.

58. A compound according to claim 57 that is (R)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.

59. A compound according to claim 57 that is (S)-8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
60. A compound according to claim 1 that is 8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
61. A compound according to claim 60 that is (R)-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
62. A compound according to claim 60 that is (S)-8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
63. A compound according to claim 1 that is 8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
64. A compound according to claim 63 that is (R)-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
65. A compound according to claim 63 that is (S)-8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
66. A compound according to claim 1 that is 8-trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.

67. A compound according to claim 66 that is (R)-8-trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
68. A compound according to claim 66 that is (S)-8-trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
69. A compound according to claim 1 that is 8-trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
70. A compound according to claim 69 that is (R)-8-trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
71. A compound according to claim 69 that is (S)-8-trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
72. A compound according to claim 1 that is 8-chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
73. A compound according to claim 72 that is (R)-8-chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
74. A compound according to claim 72 that is (S)-8-chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.


75. A compound according to claim 1 that is 8-bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
76. A compound according to claim 75 that is (R)-8-bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
77. A compound according to claim 75 that is (S)-8-bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
78. A compound according to claim 1 that is 8-iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
79. A compound according to claim 78 that is (R)-8-iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
80. A compound according to claim 78 that is (S)-8-iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
81. A compound according to claim 1 that is 7,8-dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
82. A compound according to claim 81 that is (R)-7,8-dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
83. A compound according to claim 81 that is (S)-7,8-dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.



84. A compound according to claim 1 that is 7,8-dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
85. A compound according to claim 84 that is (R)-7,8-dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
86. A compound according to claim 84 that is (S)-7,8-dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
87. A compound according to claim 1 that is 8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
88. A compound according to claim 87 that is (R)-8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
89. A compound according to claim 87 that is (S)-8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
90. A compound according to claim 1 that is 8-chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
91. A compound according to claim 90 that is (R)-8-chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.

92. A compound according to claim 90 that is (S)-8-chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
93. A composition comprising a mixture of *R* and *S* enantiomers of 8-chloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
94. A composition comprising a mixture of *R* and *S* enantiomers of 8-bromo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
95. A composition comprising a mixture of *R* and *S* enantiomers of 8-iodo-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
96. A composition comprising a mixture of *R* and *S* enantiomers of 8-trifluoromethyl-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
97. A composition comprising a mixture of *R* and *S* enantiomers of 8-trifluoromethyl-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
98. A composition comprising a mixture of *R* and *S* enantiomers of 8-chloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.

99. A composition comprising a mixture of *R* and *S* enantiomers of 8-bromo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
100. A composition comprising a mixture of *R* and *S* enantiomers of 8-iodo-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
101. A composition comprising a mixture of *R* and *S* enantiomers of 7,8-dichloro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
102. A composition comprising a mixture of *R* and *S* enantiomers of 7,8-dichloro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
103. A composition comprising a mixture of *R* and *S* enantiomers of 8-chloro-7-fluoro-1-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
104. A composition comprising a mixture of *R* and *S* enantiomers of 8-chloro-7-fluoro-1-ethyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine or a pharmaceutically acceptable salt, solvate or hydrate thereof.
105. A compound of Formula (I):



wherein:

$R_1$  is H or straight-chained or branched  $C_{1-8}$  alkyl;

$R_2$  is methyl, ethyl,  $-\text{CH}_2\text{-O-C}_{1-8}$  alkyl,  $-\text{C(=O)-O-C}_{1-8}$  alkyl,  $-\text{C(=O)-NH-C}_{1-8}$  alkyl, or  $\text{CH}_2\text{OH}$ ;

$R_{2a}$  is H;

or  $R_2$  and  $R_{2a}$  together form  $-\text{CH}_2\text{-CH}_2-$ ;

$R_3$  is H, halogen, perhaloalkyl, CN,  $\text{OR}_5$ ,  $\text{SR}_5$ ,  $\text{NHR}_5$ ,  $\text{N(R}_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $\text{C}_{1-8}$  alkyl, halogen, perhalo alkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $\text{C}_{1-8}$  alkyl;

$R_4$  is halogen, perhaloalkyl, CN,  $\text{OR}_5$ ,  $\text{SR}_5$ ,  $\text{NHR}_5$ ,  $\text{N(R}_5)_2$ , OH, aryl, or heteroaryl, wherein said aryl can be optionally substituted with up to two substituents selected from  $\text{C}_{1-8}$  alkyl, halogen, perhalo alkyl, and alkoxy, and said heteroaryl can be optionally substituted with up to two substituents selected from halogen and  $\text{C}_{1-8}$  alkyl;

or  $R_3$  and  $R_4$  together with the atoms to which they are attached can form a 5- or 6-member heterocyclic ring having one O atom;

each  $R_5$  is independently  $\text{C}_{1-8}$  alkyl,  $\text{C}_{1-8}$  alkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl or perhaloalkyl, or allyl; and

$R_6$  is H or  $\text{C}_{1-8}$  alkyl; or a pharmaceutically acceptable salt, solvate or hydrate thereof  
provided that:

(A) if  $R_2$  is methyl and  $R_1$  and  $R_3$  are both H, then  $R_4$  is not thiazole, substituted thiazole or a thiazole derivative;

(B) if  $R_6$  is other than H, then neither  $R_3$  nor  $R_4$  can be H; and

(E) if  $R_3$  is OH, and  $R_1$  is methyl then  $R_2$  cannot be cyclopentyl,  $-\text{CH}_2\text{-cyclohexyl}$ , cyclopropylmethyl, or cyclohexyl.

106. A method of prophylaxis or treatment of depression or anxiety comprising administering to a patient in need of such treatment or prophylaxis a pharmaceutically effective amount of a compound according to any one of claims 1, 42, 51, 52, 53, and 105.

107. A method of prophylaxis or treatment of a disorder related to  $5HT_{2C}$  receptor activity wherein said disorder is selected from disorders of the central nervous system; damage to the central nervous system; and gastrointestinal disorders, comprising administering to a patient in need of such prophylaxis or treatment a pharmaceutically effective amount of a compound according to any one of claims 1, 42, 51, 52, 53, and 105.

108. The method of claim 107 wherein the disorders of the central nervous system are selected from the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.

109. A method of prophylaxis or treatment of diabetes insipidus, obesity-related cardiovascular disorders, or sleep apnea comprising administering to a patient in need of such prophylaxis or treatment an effective dose of a compound according to any one of claims 1, 42, 51, 52, 53, and 105.

110. A compound according to any one of claims 1, 42, 51, 52, 53, and 105 for use in a method of treatment of the human or animal body by therapy.

111. A compound according to any one of claims 1, 42, 51, 52, 53, and 105 for use in a method of prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus; and sleep apnea.

112. The compound of claim 111 wherein disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.

113. The compound of claim 111 wherein the disorder of the central nervous system is obesity.

114. A compound according to any one of claims 1, 42, 51, 52, 53, and 105 for the manufacture of a medicament for use in the prophylaxis or treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus; and sleep apnea.

115. The compound of claim 114 wherein the disorders of the central nervous system are selected the group consisting of depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioral disorders, behavioral disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.

116. The compound of claim 114 wherein the disorder of the central nervous system is obesity.